

Predictors of Depressive Symptoms Among Hispanic Women in South Florida

By: [Amber L. Vermeesch](#), Rosa M. Gonzales-Guarda, Rosemary Hall, Brian E. McCabe, Rosina Cianelli, Nilda P. Peragallo

Vermeesch, A., Gonzalez-Guarda, R., Hall, R. F., McCabe, B., Cianelli, R. & Peragallo, N. (2013). Predictors of Depressive Symptoms among Community Dwelling Adult Hispanic Women. *Western Journal of Nursing Research*, 35(10), 1325-1338. doi: 10.1177/0193945913496152. NIHMSID: NIHMS535785

*****This version © The Authors, 2013. This is not the final version. This article has been published in the *Western Journal of Nursing Research*, published by SAGE Publications. Reprinted with permission. Figures and/or pictures may be missing from this version of the document. The version of record is available at <http://dx.doi.org/10.1177/0193945913496152>, © The Authors.*****



This work is licensed under [a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](#).

Abstract:

U.S. Hispanics, especially women, experience a disproportionate amount of disease burden for depression. This disparity among Hispanic women necessitates examination of factors associated with depression. The objective of this study was to use an adaptation of the Stress Process Model to test whether self-esteem mediated the relationship between Hispanic stress and depressive symptoms. Data for this secondary analysis were from a previous randomized-control HIV prevention trial. Participants were 548 Hispanic women (19-52 years). Data collection measures included the Center for Epidemiological Studies–Depression Scale, Rosenberg Self-Esteem Scale, and Hispanic Stress Scale. The bootstrap method in Mplus 6 was used to test mediation. Results indicated that self-esteem was inversely related to depression, and Hispanic stress was found to be positively related to depression. Self-esteem partially mediated the relationship between stress and depression. Strategies to improve/maintain self-esteem should be considered in future interventions for Hispanic women with depression.

Keywords: Latinas | Hispanics | depression | CES-D | Hispanic stress | self-esteem

Article:

In general, U.S. Hispanics experience health disparities related to the identification, treatment, and consequences of depression (Lewis-Fernández & Kleinman, 1994). Hispanic, meaning a person of Latin/South American, or Spanish heritage, is an all-inclusive term and it must be noted that many different subgroups exist and could have considerable differences regarding their experience with depression including amount of time lived in the U.S. (Weinick, Jacobs, Stone, Ortega, & Burstin, 2004). Depression is defined as a mental disorder with symptoms such as depressed mood, loss of interest, low self-worth, disturbed sleep or appetite, low energy, and poor concentration that leads to difficulties in a person's ability to take care of himself or herself (World Health Organization,

2010). There is conflicting evidence on whether U.S. Hispanics have a higher prevalence of depression than other U.S. racial and ethnic groups (Caetano & Cunradi, 2003; Delgado et al., 2006). However, Hispanics are less likely to have access to or utilize physical and mental health care services and are less likely to have their depressive symptoms identified and treated (Schiller, 2012). The lack of culturally appropriate health care services to correctly identify depressive symptoms for Hispanic populations puts them at disproportionate risk compared with other populations for unidentified and untreated depression and its consequences. This is particularly true for Hispanic women, who are more likely to experience depressive symptoms than men (Centers for Disease Control and Prevention [CDC], 2010). The literature shows that women are twice as likely to suffer depression as compared with men and in several cases the diagnosis is often associated with somatic complaints. These types of complaints serve as barriers to establishing effective prevention strategies among these women. Also, depressed women are more accepting of their symptoms, have higher thresholds for suffering and for not seeking treatment for their symptoms (Afifi, 2007; Amin & Bentley, 2002; CDC, 2010; Ferrer, Cianelli, & Peragallo, 2004).

Depression is one of the factors that increase a woman's vulnerability to acquire HIV (Brown et al., 2006; Cheng & Chan, 2007; DiClemente et al., 2001). Klein, Elifson, and Sterk (2008) stated that the presence of depressive symptoms is an aspect that adversely affects women's attitudes toward condom use and increases HIV risk behaviors. This relationship can be associated with the fact that people with high levels of depressive symptoms give less value to their lives, grounded on the feeling of hopelessness that surrounds them, and they are less willing to participate in behaviors that protect against HIV acquisition (Cheng & Chan, 2007; Myers et al., 2002; Van der Does, 2002).

The stress process, as described by Pearlin, Menaghan, Morton, and Mullan (1981), has been used to conceptualize the interacting relationships of stress, self-esteem, and depression in the unique cultural context of Hispanic women (Berry, 1997; Stroup, Lawrence, & Trevino, 1992). The stress process is conceptualized into three domains: (a) source of stress; (b) mediators of stress, conditions that can alter the impact of the sources of stress; and (c) manifestation of stress resulting in physical and psychological symptoms (Pearlin et al., 1981). In the stress process, stress and self-esteem may be independent of depressive symptoms, and self-esteem may be a mediator of the relationship between stress and depressive symptoms. The following paragraphs describe the literature that link studies between (a) Hispanic stress and depressive symptoms, (b) self-esteem and depressive symptoms, and (c) Relationship Pathways in the Stress Process Model.

Hispanic stress describes events that are often experienced by immigrants or members of nonmajority ethnic groups, specifically for Hispanics. Hispanic stress is often defined by occupational/economic stress, marital stress, immigration stress, and family/culture stress factors. Previous studies have found that these Hispanic stress factors are related to depression (Cervantes, Padilla, & Salgado de Snyder, 1991; González, Haan, & Hinton, 2001; Posner, Stewart, Marín, & Pérez-Stable, 2001). In a study conducted by De Santis, Gonzalez-Guarda, and Vasquez (2012), it was found that depression and Hispanic stress were significantly correlated ($r = .33$). Despite evidence for the link between components of Hispanic stress and depression through the stress process model for Hispanic men, little research has been conducted to test their relationships among Hispanic women.

Self-esteem has been defined as a continuum of self-worth (Rosenberg, 1965). Self-esteem is inversely related to depressive symptoms among Hispanic women (González-Guarda, Peragallo, Vasquez, Urrutia, & Mitrani, 2009; Rosenberg, 1965). Various researchers have concluded that

self-esteem is inversely related to depressive symptoms in Hispanics (De Santis et al., 2012; González-Guarda et al., 2009). As previously mentioned in the study conducted by De Santis et al. (2012), the sample was comprised of almost exclusively immigrant adult Hispanic men living with HIV. The sample in the study conducted by González-Guarda et al. (2009) comprised adult Hispanic women, the great majority were immigrants. In both of these studies, depressive symptoms were measured by the Center for Epidemiological Studies–Depression Scale (CES-D; Radloff, 1977) and self-esteem by the Rosenberg Self-Esteem Scale (RSE; 1965). Although these studies provide support for the strong inverse relationship between self-esteem and depression among Hispanic men and women, they were limited to main effect analyses that did not explore multiple relationship pathways in which stress, self-esteem, and depression may be related.

Self-esteem is one common theoretical mechanism through which stress might influence depression (Pearlin et al., 1981). Several studies described hereinbefore have linked stressors unique to Hispanics and self-esteem to depression, but only one study was found that attempted to link these constructs using a stress process model in which self-esteem mediated the link between stress and depressive symptoms (Land & Hudson, 2004). Land and Hudson (2004) conducted a cross-sectional purposive sample of 154 Latina AIDS caregivers. The researchers found that low self-esteem predicted depression for Latinas, and that self-esteem partially mediated the relationship between stress of caregiving and depression.

Purpose

The current study was designed to expand the understanding of Hispanic stress, self-esteem, and depressive symptoms and the Stress Process Model for Hispanic women. The objectives of the current study were to (a) evaluate the relationship of Hispanic stress and self-esteem to depressive symptoms among Hispanic women and (b) examine whether self-esteem mediated the relationship between Hispanic stress and depression. If distinct predictors of depressive symptoms among Hispanic women can be identified, then health care professionals, and especially nurses who care for this population, would be better prepared to identify and provide appropriate treatment for Hispanic women with depression.

Method

Design

Participants. Baseline data (N = 548) from a randomized trial of Salud-Health, Educación—Education, Prevención—Prevention, Autocuidado—Self-Care (SEPA) were used for this study (Peragallo, Gonzalez-Guarda, McCabe, & Cianelli, 2012). SEPA was a culturally tailored HIV prevention program specifically designed for Hispanic women. Eligibility criteria for the study included identifying oneself as a female Hispanic and being sexually active within the past 3 months. Participants were recruited from (a) a community-based social service organization for Hispanics, (b) an urban Florida Department of Health site, (c) fliers posted in the community, and (d) public service messages delivered via radio and televon. Nearly three quarters (74%) of women in this sample reported their health as either good or very good in the past 3 months. They represented 22 countries of origin with the highest proportions from Colombia (34%), Cuba (13%), and Peru (8%). Of the 548 women, 343 (63%) reported they were uninsured and 369 (67%) reported being unemployed at the time of the data collection. Of the women, 22% had no children

and 57% had 1 or 2 children. Monthly family income was reported in the following categories: (a) below US\$1000 (151 women, 28%), (b) between US\$1000 and US\$1999 (224 women, 41%), (c) between US\$2000 and US\$2999 (100 women, 18%), (d) between US\$3000 and US\$3999 (43 women, 8%), and (e) greater than US\$4000 (24 women, 4%). Six women (1%) did not report their income. Table 1 presents additional characteristics of the Hispanic women in the current study.

Table 1. Characteristics, CES-D, and RSE Scores of Hispanic women in study (N = 548).

Characteristics	M	SD	Range
Age	38.65	8.54	19-52
Education	13.37	3.45	0-26
% of lifetime in United States	30	26.93	0-0.98
RSE	24.49	4.86	7-30
CES-D	17.92	11.65	2-52

Note. CES-D = Center for Epidemiological Studies of Depression Scale; RSE = Rosenberg Self-Esteem Scale.

Data collection procedures. The study was approved by the Institutional Review Boards of the University of Miami in Coral Gables, Florida and the Florida State Department of Health in Tallahassee, Florida. Data were collected through standardized, face-to-face interviews, which were conducted in the preferred language (Spanish or English) of the participant by bilingual female assessors between January 2008 and April 2009 at a local community organization and a nearby research office. All measures were available in English or Spanish. The baseline interviews took approximately 3 hr and a monetary incentive (US\$50) was given to the participants for their time, travel, and childcare expenses.

Measures

Hispanic stress was measured using the Hispanic Stress Inventory (HSI; Cervantes et al., 1991). The HSI consists of 73 items with five subscales: (a) Occupational/Economic Stress, (b) Marital Stress, (c) Family/Cultural Stress, (d) Immigration Stress, and (e) Parental Stress. Parental Stress was not administered in the current study, because not all of the women were mothers, and parenting was not a focus of this study. Cronbach's alpha for the subscales in this study ranged from .74 to .83. This instrument assesses two aspects of each subscale: frequency (referring to whether a stressful event has occurred) and appraisal (referring to the respondent's perception of emotional reactions to the stressful event). Owing to the participant's perceptions of a stressful event being more informative than how often an event occurred, for the purposes of the current study the authors chose to examine only the appraisal aspect.

Self-esteem was measured using the RSE (Rosenberg, 1965). This scale has 10 items measuring an individual's perception of self-worth with item responses on a 4-point Likert-type scale ranging from strongly agree to strongly disagree. The total score ranges from 0 to 30, with higher scores indicating greater self-esteem. Internal consistency reliability in this study was good, Cronbach's $\alpha = .84$. Self-esteem was positively skewed, resulting in a square root transformation being used for analysis.

Depressive symptoms were assessed with the CES-D (Radloff, 1977), a reliable measure of depressive symptoms that has been used as a screening tool and for research in clinical depression for the past 30 years. This 20-item questionnaire used a Likert-type response format to

measure frequency of depressive symptoms during the past 1 week. Respondents are considered to have depressive symptoms when they score ≥ 16 . English and Spanish versions have been widely used with Hispanics over the past three decades (Cervantes et al., 1991; González et al., 2001; Kim, Chiriboga, & Jang, 2009; Roberts, Vernon, & Rhoades, 1989). In the current study, the internal reliability of the CES-D was strong, Cronbach's $\alpha = .94$. Owing to the depressive symptoms variable being positively skewed, a square root transformation was used to make the distribution more normal.

Data Analysis

Structural Equation Modeling (SEM) using Mplus version 6.21 (Muthen & Muthen, 1998-2007) was used for all analyses. Full information maximum likelihood estimation allowed the inclusion of cases with missing data. Preliminary measurement modeling was used to create one Hispanic stress latent variable. The following cutoffs for model fit indicated that good was Comparative Fit Index (CFI) $> .95$ and Root Mean Square Error of Approximation (RMSEA) $< .05$; and "acceptable" was CFI $> .90$, and RMSEA $< .08$ (Musil, Jones, & Warner, 1998). First, a measurement model with a single latent factor explaining variation in the four Hispanic stress subscales was tested. Next, we tested whether Hispanic stress and/or self-esteem were related to depression in an SEM model. If both predictors were significantly related to depression, mediation was tested using the bootstrap method described by MacKinnon, Lockwood, and Williams (2004). This method indicates standard error estimates by resampling and correcting for bias in the central tendency of the model estimates, and resulted in a coefficient, 95% confidence interval, and p value associated with the product of the paths to and from the mediator.

Results

Measurement Model

Table 2. Measurement Models of Hispanic Stress Subscales.

Subscale	Model 1				Model 2				
	β	<i>B</i>	<i>SE</i>	<i>p</i>	β	<i>B</i>	<i>SE</i>	<i>p</i>	
OES	.85	6.35	0.32	<.001	.66	4.94	0.32	<.001	
MS	.66	5.22	0.36	<.001	.79	6.21	0.33	<.001	
IS	.70	8.07	0.49	<.001	.48	5.52	0.53	<.001	
FCS	.66	5.69	0.39	<.001	.79	6.86	0.37	<.001	
				$\chi^2(df = 2) = 108.22, p < .00,$ CFI = .867, RMSEA = .311					$\chi^2(df = 1) = 2.33, p < .001,$ CFI = .998, RMSEA = .049

Note. Model 2 includes covariance between immigration stress and occupational/economic stress error covariance. OES = Occupational/economic stress; MS = Marital stress; IS = Immigration stress; FCS = Family/Cultural stress; CFI = Comparative Fit Index; RMSEA = Root Mean Square Error of Approximation.

Relationships to Depression

A model with a latent Hispanic stress and observed self-esteem variables predicting depressive symptoms did not have acceptable fit, CFI = 0.90, RMSEA = 0.168. Hispanic stress was

significantly related to depressive symptoms, $B = 0.77$, $SE = 0.07$, $p < .001$. Self-esteem was significantly inversely related to depressive symptoms, $B = -1.21$, $SE = 0.11$, $p < .001$.

Mediation

The mediation model included a path between Hispanic stress and self-esteem, and the indirect path from Hispanic stress through self-esteem to depressive symptoms. The mediation path is displayed in Figure 1. The mediation model had an acceptable fit to the data, $CFI = .989$, $RMSEA = .059$. Hispanic stress was significantly inversely related to self-esteem, $B = 0.25$, $SE = 0.03$, $p < .001$, and accounted for 23% of the variation in self-esteem. Hispanic stress was significantly related to depressive symptoms, $B = 0.81$, $SE = 0.06$, $p < .001$. Self-esteem was significantly inversely related to depressive symptoms, $B = -1.02$, $SE = 0.13$, $p < .001$. Hispanic stress and self-esteem jointly explained 52% of the variance in depressive symptoms. The indirect effect from Hispanic stress through self-esteem to depressive symptoms was significant, $B = 0.16$, $SE = 0.02$, $p < .001$, 95% CI [0.19, 0.32], which indicated that self-esteem mediated the relationship between Hispanic stress and depressive symptoms.




Figure 1. Self-esteem partially mediated the relationship between latent Hispanic stress and depressive symptoms.

Note. Standardized coefficients. OES = occupational/economic stress; MS = marital stress; IS = immigration stress; FCS = family/cultural stress.

*** $p < .001$.

Discussion

This study examined the relationships between Hispanic stress, self-esteem, and depression. In this study, Hispanic stress was defined and measured as the compilation of various common factors that contribute to stressful situations among U.S. Hispanics, including occupational, economic, marital, family/cultural, and immigration stressors. Taken together, they were conceptualized from a stress process framework, to predict depressive symptoms. Findings from this study indicated that Hispanic stress was positively associated with depressive symptoms, suggesting that with higher levels of reported stress, more severe depressive symptoms were experienced by the individual. Although few studies have examined the effects of a comprehensive set of stressors associated with being Hispanic in the U.S. and having depressive symptoms, this study did find this relationship to be consistent with the Stress Process Model (Pearlin et al., 1981). The Stress

Process Model can further be used to guide research that clarifies the unique contributions of specific stressors and how these interact to contribute to psychological and physical health outcomes. Determining the effects of stressors is important to identify the most important targets for intervention (e.g., occupational vs. immigration, or occupational and immigration) that have the most potential for minimizing health disparities among Hispanics.

In this study, higher reported self-esteem was expected to mediate the psychological impact of Hispanic stress on depressive symptoms. The findings from this study are consistent with reports by other researchers working with diverse Hispanic populations in the U.S. regarding the protective effect that self-esteem has on depressive symptoms (Cianelli et al., 2013; González-Guarda et al., 2009; Iwata, Turner, & Lloyd, 2002; Rosenberg, 1965). Previous research studies included samples consisting primarily of Mexican Americans, Puerto Ricans, and Cubans. Larger numbers of women from Colombia, Cuba, and Peru were in this sample; all previously understudied subgroups which strengthen the argument that self-esteem plays an important role on psychological health outcomes across Hispanic subgroups. Although this appears to be an area where there is consistency across studies, future research is needed to be conducted to clearly describe the demographic, cultural, and social–environmental contributors to self-esteem and depression across Hispanic subgroups. These studies should examine interethnic and intracultural variations that exist across racial/ethnic groups and cultural subgroups. Women’s psychological well-being is important for society because depressive symptoms interfere with a woman’s and her family’s daily life and normal functioning. The state of depression causes distress for the woman, the family, and those who care about her. This is important considering that women are the main support and care providers for the family (Afifi, 2007; Cianelli et al., 2013). In addition, there is a need for better diagnosis and treatment of mental health needs of women, partially related to their culturally reinforced subordinate position in relation to their male partners. Reducing depression is important for women’s general health and as a factor in enabling them to engage in healthy sexual behaviors (Cianelli et al., 2012).

When considering the results of this study, it is important to note that the CES-D, a much debated measure of depressive symptoms, was used in this analysis (Cervantes et al., 1991; Crockett, Randall, Shen, Russell, & Driscoll, 2005; Kim et al., 2009; Roth, Ackerman, Okonkwo, & Burgio, 2008). The interpretations of items on either the RSE or the CES-D may differ, and therefore limit generalizability, from Hispanics who were not of Mexican and Puerto Rican descent as in the primary study samples (Pascoe, Stolfi, & Ormond, 2006). Second, the study findings indicated high levels of self-esteem. Third, all the measures used were self-reported. Further research is needed to evaluate the cultural equivalency of these measures across Hispanic subgroups, and then refine or enhance these measures to adapt for cultural equivalency, if needed. When possible, these self-report measures should be combined with diagnostic verification and biological measures such as salivary or hair cortisol to assess the stress response. Fourth, the participants were participating in a HIV prevention study. Although all the data were taken from the baseline assessment, prior to any involvement in HIV prevention activities, the fact that participants were a convenience sample and willing to participate in HIV prevention activities may limit generalizability of study findings to the general population of Hispanic women. Finally, because this study was a cross-sectional design, cause-and-effect relationships cannot be established.

To advance the field of health disparities research related to stressors experienced among Hispanic women and improve their psychological well-being, it is important to develop and formally test culturally specific interventions that address common stressors among Hispanic

women living in the U.S. experience. This study's findings indicate the effect of self-esteem on depressive symptoms which contribute to knowledge related to the stressors unique to primarily immigrant Hispanic women. This new knowledge of the relationships among Hispanic stress, self-esteem, and depressive symptoms provide insight on the contribution of self-esteem to depressive symptoms and should be used in future interventions. Through this study, it was found that depression associated with greater stress load is partially mediated by the presence of self-esteem. Interventions designed to improve self-esteem may be able to mitigate depression related to stress. Such interventions could include: family dynamics and roles and coping strategies with economical/occupational stressors. Gonzalez-Guarda and colleagues (in press) found that family support had a significant interaction with Hispanic stress in the context of the Syndemic Factor (e.g. latent variable of substance abuse, violence, risky sexual behavior, and depression), and should be considered when developing future interventions to reduce stress and increase self-esteem among Hispanic women living in the United States. In essence, strategies for Hispanics in the United States are needed to reduce stressors and to promote self-esteem. Nurses in primary public health settings are uniquely qualified in their pivotal positions as hands-on caregivers to implement tailored and culturally sensitive interventions for Hispanic women.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was funded by the National Center on Minority Health and Health Disparities (NCMHD) Grant 1P60 MD002266—Center of Excellence for Health Disparities Research: El Centro (Nilda P. Peragallo, principal investigator).

References

- Afifi M. (2007). Gender differences in mental health. *Singapore Medical Journal*, 48, 385-391.
- Amin A., Bentley M. E. (2002). The influence of gender on rural women's illness experiences and health-seeking strategies for gynaecological symptoms. *Journal of Health Management*, 4, 229-249.
- Berry J. W. (1997). Immigration, acculturation, adaptation. *Applied Psychology: An International Review*, 46, 5-68.
- Brown L. K., Tolou-Shams M., Lescano C., Houck C., Zeidman J., Pugatch D., Lourie K. J. (2006). Depressive symptoms as a predictor of sexual risk among African American adolescents and young adults. *Journal of Adolescent Health*, 39, 444.e441-444.e448.
- Caetano R., Cunradi C. (2003). Intimate partner violence and depression among Whites, Blacks, and Hispanics. *Annals of Epidemiology*, 13, 661-665.
- Centers for Disease Control and Prevention. (2010). Data 2010: The Healthy People 2010 database. Mental health and mental disorders by race and ethnicity (all). Retrieved from <http://wonder.cdc.gov/data2010/focraceg.htm>

- Cervantes R. C., Padilla A. M., Salgado de, Snyder N. (1991). The Hispanic Stress Inventory: A culturally relevant approach to psychosocial assessment. *Psychological Assessment*, 3, 438-447.
- Cheng S., Chan A. (2007). Withdrawal, apathy and lack of vigor in late life depression: Factorial validity and relationship to diagnosis. *Aging & Mental Health*, 11, 532-537.
- Cianelli R., Ferrer L., Norr F. K., Miner S., Irrarrazabal L., Bernales M., McElmurry B. (2012). Mano a Mano-Mujer: An effective HIV prevention intervention for Chilean women. *Health Care for Women International*, 33, 321-341.
- Cianelli R., Lara L., Villegas N., Bernales M., Ferrer L., Kaelber L., Peragallo N. (2013). Impact of mano a mano-mujer, an HIV prevention intervention, on depressive symptoms among Chilean women. *Journal of Psychiatric and Mental Health Nursing*, 20, 263-272.
- Crockett L. J., Randall B. A., Shen Y., Russell S. T., Driscoll A. K. (2005). Measurement equivalence of the center for epidemiological studies depression scale for Latino and Anglo adolescents: A national study. *Journal of Consulting and Clinical Psychology*, 73, 47-58.
- Delgado P. L., Alegria M., Canive J. M., Diaz E., Escobar J. I., Kopelowicz A., Vega W. A. (2006). Depression and access to treatment among U.S. Hispanics: Review of the literature and recommendations for policy and research. *Focus*, 4, 38-47.
- De Santis J. P., Gonzalez-Guarda R., Vasquez E. P. (2012). Psychosocial and cultural correlates of depression among Hispanic men with HIV infection: A pilot study. *Journal of Psychiatric and Mental Health Nursing*, 19, 860-869.
- DiClemente R., Wingood G. M., Crosby R. A., Sionean C., Brown L. K., Rothbaum B., Davies S. (2001). A prospective study of psychological distress and sexual risk behavior among Black adolescent females. *Pediatrics*, 108, 1-6.
- Ferrer L., Cianelli R., Peragallo N. (2004). Domestic violence, and relationship with depression and level of self-esteem in Mexican and Puertorican women. *Horizonte de Enfermería*, 15, 23-36.
- González H. M., Haan M. N., Hinton L. (2001). Acculturation and the prevalence of depression in older Mexican Americans: Baseline results of the Sacramento area Latino study on aging. *Journal of the American Geriatrics Society*, 49, 948-953.
- Gonzalez-Guarda R. M., McCabe B., Vermeesch A. L., Cianelli R., Florom-Smith A. L., Peragallo N. (in press). Cultural phenomena and the syndemic factor: Substance abuse, violence, HIV and depression among Hispanic women. *Annals of Anthropological Practice*.
- González-Guarda R. M., Peragallo N., Vasquez E. P., Urrutia M. T., Mitrani V. B. (2009). Intimate partner violence, depression, and resource availability among a community sample of Hispanic women. *Issues in Mental Health Nursing*, 30, 227-236.
- Iwata N., Turner R. J., Lloyd D. A. (2002). Race/ethnicity and depressive symptoms in community-dwelling young adults: A differential item functioning analysis. *Psychiatry Research*, 110, 281-289.
- Kim G., Chiriboga D. A., Jang Y. (2009). Cultural equivalence in depressive symptoms in older White, Black, and Mexican-American adults. *Journal of the American Geriatrics Society*, 57, 790-796.

- Klein H., Elifson K., Sterk C. (2008). Depression and HIV risk behavior practices among “at risk” women. *Women & Health*, 48, 167-188.
- Land H., Hudson S. (2004). Stress, coping, and depressive symptomatology in Latina and Anglo Aids caregivers. *Psychology & Health*, 19, 643-666.
- Lewis-Fernández R., Kleinman A. (1994). Culture, personality, and psychopathology. *Journal of Abnormal Psychology*, 1, 67-71.
- MacKinnon D. P., Lockwood C. M., Williams J. (2004). Confidence limits for the indirect effect: Distribution of the product and resampling methods. *Multivariate Behavioral Research*, 39, 99-128.
- Musil C. M., Jones S. L., Warner C. D. (1998). Structural equation modeling and its relationship to multiple regression and factor analysis. *Research in Nursing & Health*, 21, 271-281.
- Muthen L., Muthen B. (1998-2007). *Mplus user’s guide* (5th ed.). Los Angeles, CA: Author.
- Myers H. F., Lesser I., Rodriguez N., Mira C. B., Hwang W.-C., Camp C., Wohl M. (2002). Ethnic differences in clinical presentation of depression in adult women. *Cultural Diversity and Ethnic Minority Psychology*, 8, 138-156.
- Pascoe J. M., Stolfi A., Ormond M. B. (2006). Correlates of mothers’ persistent depressive symptoms: A national study. *Journal of Pediatric Healthcare*, 20, 261-269.
- Pearlin L. I., Menaghan E. G., Morton A. L., Mullan J. T. (1981). The stress process. *Journal of Health and Social Behavior*, 22, 337-356.
- Peragallo N., Gonzalez-Guarda R., McCabe B., Cianelli R. (2012). The Efficacy of an HIV risk reduction intervention for Hispanic women. *AIDS and Behavior*, 16, 1316-1326.
- Posner S. F., Stewart A. L., Marín G., Pérez-Stable E. J. (2001). Factor variability of the center for epidemiological studies depression scale (CES-D) among urban Latinos. *Ethnicity & Health*, 6, 137-144.
- Radloff L. S. (1977). The CES-D Scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1, 385-401.
- Roberts R. E., Vernon S. W., Rhoades H. M. (1989). Effects of language and ethnic status on reliability and validity of the Center for Epidemiologic Studies–Depression Scale with psychiatric patients. *Journal of Nervous and Mental Disease*, 177, 581-592.
- Rosenberg M. (1965). *Society and the adolescent self-image*. Princeton, NJ: Princeton University Press.
- Roth D. L., Ackerman M. L., Okonkwo O. C., Burgio L. D. (2008). The four-factor model of depressive symptoms in dementia caregivers: A structural equation model of ethnic differences. *Psychology and Aging*, 23, 567-576.
- Schiller J. S. (2012). Summary health statistics for U.S. adults: National Health Interview Survey, 2010. *Vital and Health Statistics*, 10(252), 1-207.
- Stroup B. C. A., Lawrence R. H., Trevino F. M. (1992). CES-D factor structure among Mexican American and Puerto Rican women from single- and couple-headed households. *Hispanic Journal of Behavioral Sciences*, 14, 310-326.

Van der Does W. (2002). Cognitive reactivity to sad mood: Structure and validity of a new measure. *Behaviour Research and Therapy*, 40, 105-119.

Weinick R. M., Jacobs E. A., Stone L. C., Ortega A. N., Burstin H. (2004). Hispanic healthcare disparities: Challenging the myth of a monolithic Hispanic population. *Medical Care*, 42, 313-320.

World Health Organization. (2010). Depression. Retrieved from http://www.who.int/mental_health/management/depression/definition/en/