

***Lindgomyces angustiascus*, (Lindgomycetaceae, Pleosporales, Dothideomycetes), a new lignicolous species from freshwater habitats in the USA**

By: Huzefa A. Raja, [Nicholas H. Oberlies](#), Tamam El-Elimat, Andrew N. Miller, Steven E. Zelski, Carol A. Shearer

Raja, H. A., Oberlies, N. H., El-Elimat, T., Miller, A. N., Zelski, S. E., & Shearer, C. A. (2013). *Lindgomyces angustiascus*, (Lindgomycetaceae, Pleosporales, Dothideomycetes), a new lignicolous species from freshwater habitats in the USA. *Mycoscience*, 54(5), 353-361. doi:10.1016/j.myc.2012.12.004

Made available courtesy of Mycological Society of Japan & Elsevier:

<http://dx.doi.org/10.1016/j.myc.2012.12.004>

This is the author's version of a work that was accepted for publication in *Mycoscience*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Mycoscience*, [54, 5, (2013)] DOI: 10.1016/j.myc.2012.12.004

Abstract:

In this paper, we describe and illustrate a new species, *Lindgomyces angustiascus*, from submerged wood in lotic and lentic habitats from Florida, North Carolina and Wisconsin, USA. The new species is characterized by black, partially immersed, flattened, globose ascomata; fissitunicate, long, slender, obclavate asci; and one-septate, hyaline, fusiform ascospores with bipolar appendages, each covered with a gelatinous cap. Maximum Likelihood and Bayesian analyses of partial 18S nrDNA and 28S nrDNA, as well as the entire nuclear ribosomal internal transcribed spacer (ITS1, 5.8S and ITS2) region support the placement and establishment of this new species in the Lindgomycetaceae, Pleosporales, Dothideomycetes. Chemical analysis of the organic extract of *L. angustiascus* revealed the presence of 6*E*,9*E*-octadecadienoic acid and ergosterol peroxide as major secondary metabolites.

Keywords: Aquatic fungi | Phylogenetics | Saprobe | Secondary metabolites | Systematics

Article:

1. Introduction

Lindgomyces K. Hiray., Kaz. Tanaka & Shearer (Lindgomycetaceae, Pleosporales, Dothideomycetes) is a recently established freshwater ascomycete genus, which currently includes six species, *L. ingoldianus* (Shearer & K.D. Hyde) K. Hiray., Kaz. Tanaka & Shearer (type species), *L. apiculatus* K. Hiray. & Kaz. Tanaka, *L. breviappendiculatus* (Kaz. Tanaka, Sat.

Hatak. & Y. Harada) K. Hiray. & Kaz. Tanaka, *L.cinctosporae* Raja, A.N. Mill. & Shearer, *L. lemonweirensis* Raja, A.N. Mill. & Shearer, and *L. rotundatus* K. Hiray. & Kaz. Tanaka (Shearer et al. 2009; Hirayama et al. 2010; Raja et al. 2011a). It is characterized by globose to subglobose ascomata; fissitunicate, cylindrical to clavate asci that are rounded at the apex; and one-septate, hyaline ascospores with a gelatinous sheath, which extends to form bipolar mucilaginous appendages (Hirayama et al. 2010).

As part of our recent investigations of freshwater ascomycetes in North Carolina, we discovered an undescribed species of *Lindgomyces* (G202-1) occurring on submerged wood in a lake from the Piedmont Plateau. The same species was previously recorded as an undescribed *Massarina* sp. (F60-1, 2) during a study on the latitudinal, substrate, and habitat distribution patterns of freshwater ascomycetes on submerged wood in Florida (Raja et al. 2009), as well as on submerged wood in a river in Wisconsin (as A640-1). Based on evaluation of morphological characters and combined phylogenetic analysis of partial 18S nrDNA (small subunit-SSU) and 28S nrDNA (large subunit-LSU), as well as the entire nuclear ribosomal internal transcribed spacer (ITS) region, we describe and illustrate a novel species discovered in three different freshwater habitats in the USA. In addition, as part of ongoing investigations of the chemical mycology of freshwater fungi, we screened one of the isolates of the new species (G202) for secondary metabolites production.

2. Materials and methods

2.1. Sample collection and morphological examination

Collection methodology and morphological examination of samples followed outlined methods (Shearer et al. 2004; Raja et al. 2009). Specimens were deposited in the University of Illinois Herbarium (ILL). The fungal cultures are maintained at the University of Illinois, Plant Biology Fungal Culture Collection and at the University of North Carolina at Greensboro, Department of Chemistry and Biochemistry Fungal Culture Collection.

2.2. DNA extraction and PCR amplification

DNA extraction and PCR amplification of SSU and LSU nrDNA regions were performed following published procedures (Shearer et al. 2009; Hirayama et al. 2010). For amplification of ITS, a combination of ITS1F/ITS1 and ITS4 primers were used (White et al. 1990; Gardes and Bruns 1993) using established thermocycler parameters (Promputtha and Miller 2010).

2.3. Taxon sampling and phylogenetic analyses

We compiled two datasets for phylogenetic analyses: (a) a combined SSU and LSU dataset that consisted of 66 taxa, which represented a number of families currently included in the Pleosporales, Dothideomycetes (Schoch et al. 2009; Lumbsch and Huhndorf 2010; Zhang et al. 2012); and (b) an ITS dataset, which consisted of 18 taxa including three strains of the new

taxon, two strains of *L. apiculatus*, two strains of *L. breviappendiculatus*, one strain of *L. cinctosporae*, two strains of *L. ingoldianus*, one strain of *Lindgomyces* sp., two strains of *L. lemonweirensis*, and four strains of *L. rotundatus*. *Massariosphaeria typhicola* (P. Karst.) Leuchtm., which is a sister species of *Lindgomyces* (Hirayama et al. 2010; Raja et al. 2011a), was used as the outgroup taxon in the ITS phylogeny, while two taxa in the Dothideales were defined as outgroup for the combined SSU and LSU alignment. Sequences of a number of taxa were obtained from previous studies on Lindgomycetaceae (Shearer et al. 2009; Hirayama et al. 2010; Raja et al. 2011a). In addition, we also included taxa from recently introduced freshwater fungal families in the Pleosporales, such as Amniculicolaceae and Lentitheciaceae (Schoch et al. 2009; Shearer et al. 2009; Zhang et al. 2009a, b). Table 1 lists all taxa used in this study along with their GenBank numbers.

Table 1. Species used in this study.

Species	Voucher info ^b	GenBank accession numbers ^a		
		nucSSU rDNA	nuc ITS	nucLSU rDNA
<i>Amniculicola immersa</i>	CBS 123083	GU456295	–	FJ795498
<i>Amniculicola lignicola</i>	CBS 123094	EF493863	–	EF493861
<i>Amniculicola parva</i>	CBS 123092	GU296134	–	FJ795497
<i>Anguillospora longissima</i>	CS869-1D	GU266222	–	GU266240
<i>Byssothecium circinans</i>	CBS 675.92	AY016339	–	AY016357
<i>Bimuria nova-zelandica</i>	CBS 107.79	AY016338	–	AY016356
<i>Cochliobolus heterostrophus</i>	CBS 134.39	AY544727	–	AY544645
<i>Cucurbitaria elongata</i>	CBS 171.55	U42482	–	DQ678061
<i>Delitschia didyma</i>	UME 31411	AF242264	–	DQ384090
<i>Delitschia winteri</i>	CBS 225.62	DQ678026	–	DQ678077
<i>Didymella cucurbitacearum</i>	IMI 373225	AY293779	–	AY293792
<i>Didymella exigua</i>	CBS 183.55	EU754056	–	EU754155
<i>Dothidea insculpta</i>	CBS 189.58	DQ247810	–	DQ247802
<i>Dothidea sambuci</i>	DAOM 231303	AY544722	–	NG_027611.1
<i>Herpotrichia juniperi</i>	CBS 468.64	U42483	–	DQ384093
<i>Lentithecium aquaticum</i>	CBS 123099	FJ795477	–	FJ795434
<i>Lentithecium fluviatile</i>	CBS 123090	FJ795492	–	FJ795450
<i>Lepidosphaeria nicotiae</i>	CBS 559.71	DQ384068	–	DQ384068
<i>Leptosphaeria biglobosa</i>	CBS 532.66	EU754090	–	EU754189
<i>Letendraea helminthicola</i>	CBS 884.85	AY016345	–	AY016362
<i>Lindgomyces angustiascus</i>	A640-1a Type	JX508280	JX508281	JX508279

<i>Lindgomyces angustiascus</i>	A640-1b	–	JX508282	–
<i>Lindgomyces angustiascus</i>	F60-1	JX508284		JX508283
<i>Lindgomyces angustiascus</i>	G202-1a	JX508287	JX508286	JX508285
<i>Lindgomyces apiculatus</i>	JCM 13091/MAFF 239601 TYPE	JF419886	JF419892	JF419884
<i>Lindgomyces apiculatus</i>	JCM 13092/MAFF 239602	JF419887	JF419893	JF419885
<i>Lindgomyces breviappendiculatus</i>	JCM 12702/MAFF 239291	AB521734	JF419896	AB521749
<i>Lindgomyces breviappendiculatus</i>	JCM 12701/MAFF 239292 Type	AB521733	JF419897	AB521748
<i>Lindgomyces cinctosporae</i>	R56-1 Type	AB522430	JF419905	AB522431
<i>Lindgomyces cinctosporae</i>	R56-3	GU266238	–	GU266245
<i>Lindgomyces ingoldianus</i>	ATCC 200398 Type	AB521719	JF419898	AB521736
<i>Lindgomyces ingoldianus</i>	JCM 16479/NBRC 106126	AB521720	JF419899	AB521737
<i>Lindgomyces</i> sp.	JCM 16480/NBRC 106130	AB521721	JF419900	AB521738
<i>Lindgomyces lemonweirensis</i>	A632-1a Type	JF419890	JF419894	JF419888
<i>Lindgomyces lemonweirensis</i>	A632-1b	JF419891	JF419895	JF419889
<i>Lindgomyces rotundatus</i>	JCM 16481/MAFF 239473 Type	AB521722	JF419901	AB521739
<i>Lindgomyces rotundatus</i>	JCM 16482/NBRC106127	AB521723	JF419902	AB521740
<i>Lindgomyces rotundatus</i>	JCM 16483/NBRC 106128	AB521724	JF419903	AB521741
<i>Lindgomyces rotundatus</i>	JCM 16484/NBRC 106129	AB521725	JF419904	AB521742
<i>Lolia aquatica</i>	–	–	–	HM367732
<i>Lophiostoma heterosporum</i>	–	AY016345	–	AY016369
<i>Lophiostoma macrostomum</i>	JCM 13545	AB521731	–	AB433273
<i>Lophiostoma macrostomum</i>	JCM 13546/MAFF 239447	AB521732	–	AB433274
<i>Massaria platani</i>	CBS 221.37	DQ678013	–	DQ678065
<i>Massarina eburnea</i>	JCM 14422	AB521718	–	AB521735
<i>Massariosphaeria</i>	MAFF 239218	AB521729	–	AB521746

<i>typhicola</i>				
<i>Massariosphaeria typhicola</i>	MAFF 239219	AB521730	JF419906	AB521747
<i>Montagnula opulenta</i>	–	AF164370	–	DQ678086
<i>Neotestudina rosatii</i>	CBS 690.82	DQ384069	–	DQ384107
<i>Neottiosporina paspali</i>	CBS 331.37	EU754073	–	EU754172
<i>Ophiosphaerella herpotricha</i>	CBS 620.86	DQ678010	–	DQ678062
<i>Phaeosphaeria avenaria</i>	AFTOL-ID 280	AY544725	–	AY544684
<i>Phaeodothis winteri</i>	CBS 182.58	DQ678021	–	DQ678073
<i>Phoma herbarum</i>	CBS 615.75	EU754087	–	EU751486
<i>Pleospora herbarum</i>	CBS 714.68	DQ767648	–	DQ678049
<i>Pleomassaria siparia</i>	CBS 279.74	DQ678027	–	AY004341
<i>Preussia terricola</i>	DAOM 230091	AY544726	–	AY544686
<i>Setomelanomma holmii</i>	CBS 110217	AF525677	–	AF525678
<i>Setosphaeria monoceras</i>	CBS 154.26	AY016352	–	AY016368
<i>Sporormia lignicola</i>	CBS 264.69	U42478	–	DQ384098
<i>Tingoldiagio graminicola</i>	JCM 16485/NBRC 106131 Type	AB521726	–	AB521743
<i>Tingoldiagio graminicola</i>	MAFF 239472	AB521727	–	AB521744
<i>Tingoldiagio graminicola</i>	JCM 16486/NBRC 106132	AB521728	–	AB521745
<i>Trematosphaeria pertusa</i>	CBS 400. 97	DQ678020	–	DQ678072
<i>Ulospora bilgramii</i>	CBS 110020	DQ384071	–	DQ384108
<i>Verruculina enalia</i>	CBS 304.66	DQ678028	–	AY016363
<i>Zopfia rhizophila</i>	CBS 270.26	L76622	–	DQ384104

a Number in Bold indicates newly obtained sequences in this study.

b CBS, Centraalbureau voor Schimmelcultures; A, Carol Shearer; IMI, International Mycological Institute; UME, Umeå University, Sweden; DAOM, Agriculture and Agri-Food Canada National Mycological Herbarium; JCM, Japan Collection of Microorganisms; MAFF, National Institute of Agrobiological Sciences, Japan; NBRC, National Biological Resource Center, Japan; R, Raja Freshwater Ascomycetes; F, Florida Freshwater Ascomycetes A, Carol Shearer, Ascomycetes; AFTOL, Assembling the Fungal Tree of Life; ATCC, American Type Culture Collection; G, University of North Carolina at Greensboro, (UNCG), Department of Chemistry and Biochemistry Culture Collection.

Each of the datasets was aligned initially using the multiple alignment program MUSCLE[®] (Edgar 2004) set to default parameters as implemented in the program Seaview v. 4.1 (Gouy et al. 2010). Alignments were then optimized by visual examination and corrected manually using MacClade v. 4.08 (Maddison and Maddison 2000). After the datasets were aligned, ambiguous regions, gaps and introns were excluded from the final alignment using the default parameters in the program Gblocks (Castresana 2000; Talavera and Castresana 2007).

Nucleotides from the 5' and 3' ends were also removed in both the datasets due to missing characters in most taxa.

Maximum Likelihood (ML) analyses were performed on both datasets. JModeltest2 was used (Darriba et al. 2012) (with 1624 possible evolutionary models) based on the implementation of the Akaike Information Criterion (AIC) to obtain the best-fit model of nucleotide evolution. Conflict between the individual SSU and LSU datasets was assessed using the methods outlined in Lutzoni et al. (2004) and Raja et al. (2011b). Based on results of this study as well as previous phylogenetic analyses (Shearer et al. 2009; Hirayama et al. 2010; Raja et al. 2011a), no significant conflicts between separate SSU and LSU trees were found; therefore, the two datasets were concatenated to run the ML analyses. For the combined SSU and LSU dataset, the GTR + I + G model (Rodríguez et al. 1990) was selected. For the ITS dataset, the TIM3ef + G model was selected by AIC in jModeltest2.

For both datasets, ML was performed with 1000 ML bootstrap (BS) replicates with a combined Nearest Neighbor Interchange (NNI) and Subtree Pruning and Regrafting (SPR) tree search option in effect using PHYML (Guindon and Gascuel 2003). Additional ML analyses were performed using RAxML v. 7.0.4 (Stamatakis et al. 2008) on both the combined SSU and LSU dataset as well as on the ITS dataset; these analyses were run on the CIPRES Portal v. 2.0 (Miller et al. 2010) with the default rapid hill-climbing algorithm and GTR model employing 1000 fast BS searches. Clades with a bootstrap value (BSV) $\geq 70\%$ were considered significant and strongly supported (Hillis and Bull 1993).

Bayesian analyses were performed to assess nodal support on the combined SSU and LSU dataset as well as separately on the ITS dataset using MrBayes v 3.12 (Huelsenbeck and Ronquist 2001, 2005) implementing the above models. These analyses were run on the CIPRES Portal v. 2.0 (Miller et al. 2010). Constant characters were included and 100 million generations with trees sampled every 100th generation were run resulting in 100,000 total trees for each of the combined SSU + LSU and ITS datasets. Based on information on prior runs, the first 10,000 trees extended beyond the burn-in phase in each analysis, so these were discarded and the remaining 90,000 trees were used to calculate the Bayesian posterior probability (BPP). The consensus trees were generated in PAUP 4.0b10 (Swofford 2002). The Bayesian analyses were run twice starting from a different random tree each time to ensure that trees from the same tree space were sampled.

2.4. Chemical analysis

2.4.1. Fermentation, extraction and isolation

A fresh culture of G202 was grown on 2% malt extract agar (MEA) (20 g malt extract, 15 g agar, 1000 ml sterile distilled water). After 14–21 d, a piece of agar culture was transferred to a medium containing yeast extract, soy peptone, and dextrose (YESD; 20 g soy peptone, 20 g dextrose, 10 g yeast extract, 1000 ml sterile distilled water). After incubation (7–14 d) at 22 °C

with agitation, the culture was used to inoculate 50 ml of a rice medium, prepared using 25 g of rice and 35 ml of H₂O in a 250 ml Erlenmeyer flask. This was incubated at 22 °C until the culture showed good growth (approximately 14 d). To the solid fermentation culture of G202, 150 ml of 1:1 MeOH–CHCl₃ was added. The culture was chopped with a spatula and shaken overnight (~16 h) at ~100 rpm at room temperature. The sample was vacuum filtered, and the remaining residues were washed with small volumes of 1:1 MeOH–CHCl₃. To the filtrate, 90 ml CHCl₃ and 150 ml H₂O were added; the mixture was stirred for 1/2 h and then transferred into a separatory funnel. The bottom layer was drawn off and evaporated to dryness. The dried organic extract was re-constituted in 50 ml of 1:1 MeOH–CH₃CN and 50 ml of hexane. The biphasic solution was then mixed vigorously in a separatory funnel. The MeOH–CH₃CN layer was drawn off and evaporated to dryness under vacuum. The defatted material (127.1 mg) was dissolved in a mixture of CHCl₃–MeOH, adsorbed onto Celite 545, and fractionated via a Teledyne ISCO CombiFlash Rf using a 4 g Silica column and a gradient solvent system of hexane–CHCl₃–MeOH at an 18 ml/min flow rate and 68.1 column volumes over 18.2 min to afford four fractions. Fraction 2 was eluted with 100% CHCl₃ (~13 mg) and was subjected to semipreparative HPLC purification over a Gemini-NX C18 (5 μm; 250 × 10 mm; Phenomenex, Inc., Torrance, CA, USA) column using a mobile phase consisting of CH₃CN–H₂O (acidified with 0.1% formic acid) starting with 90:10 then increasing linearly to 100% CH₃CN within 15 min at a flow rate of 4.7 ml/min to yield five sub-fractions. Sub-fraction 1 yielded compounds **1** (2.97 mg) and **2** (2.49 mg), which eluted at ~10.0 and 18.0 min, respectively.

Identification—The structures of compounds **1** and **2** were elucidated using high-resolution mass spectrometry (HRMS) and Nuclear Magnetic Resonance (NMR) spectroscopy. The HRMS was performed on a Thermo LTQ Orbitrap XL mass spectrometer equipped with an electrospray ionization source. NMR experiments were conducted in CDCl₃ with TMS as a reference via a JEOL ECA-500, operating at 500 MHz for ¹H.

3. Results

3.1. Morphological data

Examination of specimens based on whole crush mounts of fresh material indicated that the new fungus fits well within the genus *Lindgomyces*. Characteristics of this fungus not observed in any of the six previously described species of *Lindgomyces* include long, apically narrow, obclavate asci and presence of narrow, fusiform ascospores with ephemeral bipolar appendages with gelatinous caps. The morphological data support the establishment of *L. angustiascus* as a novel species within the family Lindgomycetaceae.

3.2. Molecular data

The two partial nrDNA regions were concatenated since no significant conflicts were found between the individual SSU and LSU trees. The original combined SSU and LSU alignment comprised 66 taxa and 4747 bp positions. After removal of ambiguous regions with Gblocks and

excluding large segments of missing data from the 5' and 3' ends, the final alignment included 2312 bp. The base contents were as follows: % GC = 48.07, % A = 26.156, % C = 20.828, % G = 27.979, and % T = 25.037. PHYML analyses of the combined SSU and LSU dataset produced a single most likely tree ($-\ln L = 9954.30$) (Fig. 1). All species of *Lindgomyces* occurred in a highly supported clade within the family Lindgomycetaceae (Hirayama et al. 2010; Raja et al. 2011a) with $\geq 95\%$ BPP, 80% PHYML BS, and 80% RAxML BS. Isolates of the new species, *L. angustiascus*, formed a well-supported clade [$\geq 95\%$ BPP, 77% PHYML BS, 96% RAxML BS (Fig. 1)].




Fig. 1. Phylogram of the most likely tree ($-\ln L = 9954.30$) from a PHYML analysis of 66 taxa based on combined SSU and LSU nrDNA (2312 bp). Numbers refer to PHYML/RAxML bootstrap support values $\geq 70\%$ based on 1000 replicates. An asterix indicates significant Bayesian

posterior probabilities $\geq 95\%$. Members of the Dothideales were used as outgroup taxa. The new species is shown in bold. Classification following Lumbsch and Huhndorf (2010) is shown on the right.

The original ITS alignment consisted of 18 taxa and 1197 bp positions. After using Gblocks to remove ambiguous regions from the sequence alignment and excluding the missing data from the 5' and 3' ends, the final dataset consisted of 975 bp. The length range of the ITS dataset was 476–975 bp. The base contents were as follows: % GC = 51.802, % A = 22.024, % C = 25.147, % G = 26.654, and % T = 26.174. PHYML analyses of the ITS dataset generated a single most likely tree ($-\ln L = 3448.21$, Fig. 2). Isolates of *L.angustiascus* formed a distinct monophyletic group with $\geq 95\%$ PP, 99% PHYML BS, and 100% RAxML BS. Both isolates a and b from strain A640-1 grouped together with $\geq 95\%$ PP, 99% PHYML BS, and 99% RAxML BS. *Lindgomyces angustiascus* occurred as a sister clade to *L. lemonweirensis* with $\geq 95\%$ PP, 100% PHYML BS, and 100% RAxML BS (Fig. 2). Alignments of both datasets were deposited in TreeBASE (www.treebase.org, submission number (S13461)).




Fig. 2. Phylogram of the most likely tree ($-\ln L = 3448.21$) from a PHYML analysis of 18 taxa based on ITS nrDNA (975 bp). Support values as in Fig. 1.

The molecular phylogenetic analyses of both the combined SSU and LSU (Fig. 1) as well as the ITS phylogeny (Fig. 2) added further support to the establishment of *L. angustiascus* within the Lindgomycetaceae. The new species appeared to be a phylogenetically related sister species to *L. lemonweirensis*. This taxon found in Florida, North Carolina, and Wisconsin is therefore described and illustrated herein as a new species of *Lindgomyces*.

3.3. Chemistry data

From the organic extract of *L. angustiascus* (G202), two major compounds were isolated and identified as 6*E*,9*E*-octadecadienoic acid (**1**) and ergosterol peroxide (**2**) (Suppl. Fig. 1). The spectral data compared favorably to those reported previously (Dictionary of Natural Products, www.chemnetbase.com) and matched authentic standards.

3.4. Taxonomy

Lindgomyces angustiascus Raja, A.N. Mill. & Shearer, sp. nov. Fig. 3




Fig. 3. *Lindgomyces angustiascus*. A, ascoma in longitudinal section (F60-1); B, ascomal wall (F60-1); C, pseudoparaphyses; D, asci; E, elongating ascus showing ascospores with an apical appendage; F, narrow ascus apex; G, ascospore in water showing apical appendages and gelatinous caps (arrow); H, ascospore showing apical appendages; I, ascospores in water; J, ascospore mounted in glycerin; K–N, ascospores in water. C–N: from the HOLOTYPE. Bars: A, 50 μm ; B, D–G, I, K–N, 20 μm ; C, H, J, 10 μm .

Mycobank no.: MB 801725.

Differs from other *Lindgomyces* spp. in characteristic long, slender, obclavate asci with a narrow apex; ascospores newly discharged from the asci possess bipolar apical appendages with gelatinous caps.

Type: USA. WISCONSIN: Lemonweir River at jct. with Rt. HH south of Rt. 82, 43°46'16"N, 89°53'10"W, on submerged decorticated wood, 5 October 2009, Huzefa Raja and Andrew N. Miller, A640-1 (HOLOTYPE designated here, ILL 41201).

Etymology: From L., angustus, “narrow”, referring to the narrow ascus apex of this species compared to other species of *Lindgomyces*.

Ascomata on wood 290–395 \times 270–280 μm , black, partially immersed, scattered, flattened globose, with a broad flattened base, ostiolate, papillate, clypeate. Papillae 40–50 \times 60–90 μm , broad, conical, central; wall of papilla of darkened pseudoparenchymatic cells. Peridium ca. 20–35 μm wide, two-layered; inner layer of hyaline angular cells 10–15 \times 3–5 μm , outer layer composed of dark irregularly shaped cells. Pseudoparaphyses numerous, ca. 1–2 μm wide, septate, anastomosing above the asci. Asci 122–203 \times 18–28 μm (mean = 158 \times 25 μm , $n = 40$), fissitunicate, long, slender, obclavate, narrow and thick-walled at the apex, broader and thinner walled below, slightly curved, eight-spored, overlapping uniseriate at ascus apex, biseriate at ascus base. Ascospores 47–58 \times 9–12 μm (mean = 53 \times 11 μm , $n = 70$), ellipsoidal to fusiform, tapering at the apices, hyaline, one-septate, multiguttulate, with two to three large guttules in each cell; equipped with bipolar appendage-like structures ca. 2 μm long and initially covered with gelatinous caps; appendages ephemeral in water.

Additional specimens examined: Florida, Apalachicola National Forest, Wood Lake, 30°01'34"N, 84°33'57"W, water 30 °C, pH 8.5, on submerged decorticated wood, July 2004, *Chris Brown* and *Huzefa Raja*, F60-1; Apalachicola River at Fort Gadsden Landing, 29°56'25.87"N, 85°0'40.54"W, water 9 °C, pH 6, on submerged decorticated wood, 14 February 2006, *J.L. Crane* and *Huzefa Raja*, F60-2; North Carolina, Lake at Hagan Stone Park, 35°57'9"N, 79°44'9"W, on submerged partially decorticated wood, 27 March 2012, *Huzefa Raja*, *Mario Figueroa* and *Daniella Hayes*, G202-1.

Culture: Colonies on Potato Dextrose Agar (PDA; Difco, Detroit, MI, USA) growing slowly (20 mm diam in 3 weeks), diffuse, submerged, floccose or cottony, margin even, dark gray at the center, and white toward the periphery.

Anamorph: not observed.

Known distribution: USA (Florida, North Carolina, Wisconsin).

Comments: *Lindgomyces angustiascus* can be distinguished from other species in the genus by the characteristic long, slender, obclavate or cymbiform asci with a narrow apex. Ascospores newly discharged from the asci possess bipolar apical appendages with gelatinous caps (Fig. 3G); however, the appendages are ephemeral and dissolve away quickly in water. Thus far, this species was found occurring only on submerged woody substrates, suggesting that it may prefer lignicolous substrates.

4. Discussion

Using molecular sequence data and morphological evidence, Hirayama et al. (2010) established the family *Lindgomycetaceae* based on *Lind. ingoldianus* (Basionym: *Massarina ingoldiana*), a freshwater member of the Dothideomycetes similar to *M. eburnea* (Tul. & C. Tul.) Sacc., the type species of *Massarina*. The molecular analyses, which included sequences for *M. eburnea*, *Lophiostoma macrostomum* (Tode) Ces. & de Not., and *Lind. ingoldianus*, revealed that these taxa were distantly related in a phylogenetic tree based on a wide range of Dothideomycetes genera. Molecular phylogenetic studies using ribosomal sequence data from different lineages of Dothideomycetes belonging to freshwater, marine and terrestrial bitunicate fungi also showed that *Lindgomycetaceae* was a unique lineage among the Dothideomycetes and did not share phylogenetic affinities with *Massarina sensu stricto* or *Lophiostomataceae* (Schoch et al. 2009; Shearer et al. 2009; Raja et al. 2011a). Based on the molecular phylogenetic analysis performed in this study (Fig. 1), it is evident that *L. angustiascus* belongs in the genus *Lindgomyces*, *Lindgomycetaceae* and is not conspecific with taxa such as *Massarina* or *Lophiostoma*.

Data obtained from both molecular and morphological analyses strongly support the placement of *L.angustiascus* within *Lindgomyces*. The morphological characteristics such as long, slender, obclavate asci and narrow, pointed, fusiform ascospores with ephemeral appendages covered with gelatinous caps in *L.angustiascus* distinguishes this species from all previously recognized species within *Lindgomyces*. Molecular data obtained from combined SSU and LSU sequences and ITS data (Figs. 1 and 2) also supports observations from phenotypic data (Fig. 3).

The ascospores of *Lindgomyces angustiascus* are morphologically similar to those of *L. breviappendiculatus* (Tanaka et al. 2005; Hirayama et al. 2010) and *L. apiculatus* (Raja et al. 2011a) in having bipolar apical appendages. *Lindgomyces angustiascus*, however, differs from the latter two species in having ascospores with similar sized upper and lower cells and not

being strongly constricted at the midseptum. In addition, as soon as the ascospores of *L. angustiascus* are released in water, short, gelatinous caps separate from the ascospore apices to release ephemeral, gelatinous appendages (Fig. 3G), a character not observed in other *Lindgomyces* species thus far. A key to species of *Lindgomyces* has been published by Raja et al. (2011a).

The ITS region of the ribosomal operon has been designated as the fungal barcode marker (Schoch et al. 2012). A Blast search (Altschul et al. 1997) of the ITS sequence of isolates of *L. angustiascus* in GenBank (Benson et al. 2012) suggested that the ITS sequence of the new fungus is 97–98% similar to that of *L. lemonweirensis* (JF419894 and JF419895). However, *L. angustiascus* is morphologically distinct from *L. lemonweirensis* in both ascus and ascospore dimensions as well as the morphology of the ascus; the asci of *L. angustiascus* are long, slender and cymbiform with a narrow apex, whereas those of *L. lemonweirensis* are clavate to cymbiform and rounded at the apex (Raja et al. 2011a). In addition, the ascospores of *L. angustiascus* possess bipolar apical appendages with gelatinous caps, while an entire gelatinous sheath surrounds the ascospores of *L. lemonweirensis*. Although the 3% cutoff as a proxy for ITS based species identification works reasonably well in fungi, caution must be used when translating information from sequence data into species names (Nilsson et al. 2008). Previous taxonomic studies using ITS data for species level identification (Harrington and Rizzo 1999) showed sibling species that have recently diverged did not show pronounced differences in ITS sequences, although the taxa under investigation were morphologically distinct.

As part of recent investigations regarding the chemical mycology of freshwater fungi, we are characterizing the chemical constituents of new and unusual ascomycetes collected from various lotic and lentic habitats. A fatty acid, 6*E*,9*E*-octadecadienoic acid (**1**) and ergosterol peroxide (**2**) were the major chemical compounds isolated from strain G202 (Suppl. Fig. 1). Fatty acids have been isolated commonly from filamentous fungi (Stahl and Klug 1996; Jie et al. 1998), and their profiles might be useful in understanding intraspecific variation and chemotaxonomic profiling (Stahl and Klug 1996; Frisvad et al. 2008). Additional natural products chemistry studies on *Lindgomyces* spp. could be conducted to determine if members of Lindgomycetaceae share similar or different fatty acid profiles. Ergosterol peroxide is a steroid derivative that has been isolated from a number of different filamentous fungi including both Ascomycota (Kuo et al. 2003) and Basidiomycota (Krzyczkowski et al. 2009).

It is noteworthy that four out of the seven described species currently placed in *Lindgomyces* including *L. angustiascus* have been described or collected from submerged wood in the Lemonweir River in WI, USA. This new species was collected from three distinct geographical locations along a latitudinal gradient in the USA. Additional collections of submerged woody substrates from lotic and lentic freshwater habitats in different geographical locations will certainly expand the distribution patterns of taxa within the Lindgomycetaceae. Such broad-scale geographical surveys may further add new taxa within this family and expand


the suite of morphological and molecular characters that currently encompass this freshwater ascomycete genus.

Acknowledgments

The authors from the University of North Carolina at Greensboro were supported in part by internal research and development funds. Financial support to CAS and ANM from the National Science Foundation (NSF Grant No. 08-44722) is greatly appreciated. Any opinions, findings and conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect the views of the National Science Foundation.

Appendix A. Supplementary data

The following is the supplementary data related to this article:



Supplementary Fig. 1.

Chemical structures of compounds (1) and (2).

References

- Altschul S, Madden T, Schaffer A, Zhang JH, Zhang Z, Miller W, Lipman D, 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 17: 3389e3402.
- Benson DA, Karsch-Mizrachi I, Clark K, Lipman DJ, Ostell J, Sayers EW, 2012. GenBank. *Nucleic Acids Research* 40: 48e53.
- Castresana J, 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17: 540e552.
- Darriba D, Guillermo LT, Ramon D, Posada D, 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* 9: 8.
- Edgar RC, 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792e1797.

- Frisvad JC, Andersen B, Thrane U, 2008. The use of secondary metabolite profiling in chemotaxonomy of filamentous fungi. *Mycological Research* 112: 231e240.
- Gardes M, Bruns TD, 1993. ITS primers with enhanced specificity for basidiomycetes e application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2: 113e118.
- Gouy M, Guindon S, Gascuel O, 2010. SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Molecular Biology and Evolution* 27: 221e224.
- Guindon S, Gascuel O, 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology* 52: 696e704.
- Harrington TC, Rizzo DM, 1999. Defining species in fungi. In: Worrall JJ (ed) *Structure and dynamics of fungal populations*. Kluwer Press, Dordrecht, pp 43e71.
- Hillis DM, Bull JJ, 1993. An empirical-test of bootstrapping as a method for assessing confidence in phylogenetic analysis. *Systematic Biology* 42: 182e192.
- Hirayama K, Tanaka K, Raja HA, Miller AN, Shearer CA, 2010. A molecular phylogenetic assessment of *Massarina ingoldiana* sensu lato. *Mycologia* 102: 729e746.
- Huelsenbeck JP, Ronquist F, 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754e755.
- Huelsenbeck JP, Ronquist F, 2005. Bayesian analysis of molecular evolution using Mr. Bayes. In: Nielsen R (ed) *Statistical methods in molecular ecology*. Springer Press, pp 186e226.
- Jie MSFLK, Pasha MK, Syed-Rahmatullah MSK, 1998. Fatty acids, fatty acid analogues and their derivatives. *Natural Products Report* 15: 607e629.
- Krzyczkowski W, Malinowska E, Suchocki P, Kleps J, Olejnik M, Herold F, 2009. Isolation and quantitative determination of ergosterol peroxide in various edible mushroom species. *Food Chemistry* 113: 351e355.
- Kuo YC, Weng SC, Chou CJ, Chang TT, Tsai WJ, 2003. Activation and proliferation signals in primary human T lymphocytes inhibited by ergosterol peroxide isolated from *Cordyceps cicadae*. *British Journal of Pharmacology* 140: 895e906.
- Lumbsch HT, Huhndorf SM, 2010. Myconet volume 14. Part Two. Notes on ascomycete systematics. Nos. 4751e5113. *Fieldiana: Life and Earth Science*, NS.
- Lutzoni F, Kauff F, Cox CJ, McLaughlin D, Celio G, Dentinger C, Padamsee M, Hibbett D, James TY, Baloch E, Grube M, Reeb V, Hofstetter V, Schoch C, Arnold AE, Miadlikowska J, Spatafora J, Johnson D, Hambleton S, Crockett M, Shoemaker R, Sung GH, Lu cking R, Lumbsch T, O'Donnell K, Binder M, Diederich P, Ertz D, Gueidan C, Hansen K, Harris RC,

Hosaka K, Lim YW, Matheny B, Nishida H, Pfister D, Rogers J, Rossmann A, Schmitt I, Sipman H, Stone J, Sugiyama J, Yahr R, Vilgalys R, 2004. Assembling the fungal tree of life: progress, classification, and evolution of subcellular traits. *American Journal of Botany* 91: 1446e1480.

Maddison WP, Maddison DR, 2000. *MacClade: analysis of phylogeny and character evolution*. Sinauer, Sunderland, Massachusetts.

Miller MA, Pfeiffer W, Schwartz T, 2010. Creating the CIPRES science gateway for inference of large phylogenetic trees. Paper Presented at the Proceedings of the Gateway Computing Environments workshop (GCE), 14 Nov. 2010.

Nilsson RH, Kristiansson E, Ryberg M, Hallenberg N, Larsson KH, 2008. Intraspecific ITS variability in the kingdom fungi as expressed in the international sequence databases and its implications for molecular species identification. *Evolutionary Bioinformatics* 4: 193e201.

Promptutha I, Miller AN, 2010. Three new species of *Acanthostigma* (Tubeufiaceae, Dothideomycetes) from Great Smoky Mountains National Park. *Mycologia* 102: 574e587.

Raja HA, Schmit JP, Shearer CA, 2009. Latitudinal, habitat and substrate distribution patterns of freshwater ascomycetes in the Florida Peninsula. *Biodiversity and Conservation* 18: 419e455.

Raja HA, Tanaka K, Hirayama K, Miller AN, Shearer CA, 2011a. Freshwater ascomycetes: two new species of *Lindgomyces* (Lindgomycetaceae, Pleosporales, Dothideomycetes) from Japan and USA. *Mycologia* 103: 1421e1432.

Raja HA, Schoch CL, Hustad VP, Shearer CA, Miller AN, 2011b. Testing the phylogenetic utility of MCM7 in the Ascomycota. *Myckeys* 1: 63e94; <http://dx.doi.org/10.3897/mycokeys.1.1966>.

Rodríguez F, Oliver JL, Marin A, Medina JR, 1990. The general stochastic model of nucleotide substitutions. *Journal of Theoretical Biology* 142: 485e501.

Schoch CL, Crous PW, Groenewald JZ, Boehm EWA, Burgess TI, de Gruyter J, de Hoog GS, Dixon LJ, Grube M, Gueidan C, Harada Y, Hatakeyama S, Hirayama K, Hosoya T, Huhndorf SM, Hyde KD, Jones EBG, Kohlmeyer J, Kruijs A, Li YM, Lücking R, Lumbsch HT, Marvanová L, Mbatchou JS, McVay AH, Miller AN, Mugambi GK, Muggia L, Nelsen MP, Nelson P, Owensby CA, Phillips AJL, Phongpaichit S, Pointing SB, Pujade-Renaud V, Raja HA, Plata ER, Robbertse B, Ruibal C, Sakayaroj J, Sano T, Selbmann L, Shearer CA, Shirouzu T, Slippers B, Suetrong S, Tanaka K, Volkmann-Kohlmeyer B, Wingfield MJ, Wood AR, Woudenberg JHC, Yonezawa H, Zhang Y, Spatafora JW, 2009. A class-wide phylogenetic assessment of Dothideomycetes. *Studies in Mycology* 64: 1e15.

Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W, Fungal Barcoding C, Fungal Barcoding Consortium, 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. *Proceedings of the National Academy of Sciences of the United States of America* 109: 6241e6246.

Shearer CA, Langsam DM, Longcore JE, 2004. Fungi in freshwater habitats. In: Mueller GM, Bills GF, Foster MS (eds), Biodiversity of fungi: inventory and monitoring methods. Elsevier, Amsterdam, pp 513e531.

Shearer CA, Raja HA, Miller AN, Nelson P, Tanaka K, Hirayama K, Marvanova L, Hyde KD, Zhang Y, 2009. The molecular phylogeny of freshwater Dothideomycetes. *Studies in Mycology* 64: 145e153.

Stahl PD, Klug MJ, 1996. Characterization and differentiation of filamentous fungi based on fatty acid composition. *Applied Environmental Microbiology* 62: 4136e4146. Stamatakis A, Hoover P, Rougemont J, 2008. A rapid bootstrap algorithm for the RAxML web servers. *Systematic Biology* 57: 758e771.

Swofford DL, 2002. PAUP*: phylogenetic analysis using parsimony (* and other methods). Version 4. Sinauer Associates, Sunderland, Massachusetts.

Talavera G, Castresana J, 2007. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Systematic Biology* 56: 564e577. Tanaka K, Hatakeyama S, Harada Y, 2005. Three new freshwater ascomycetes from rivers in Akkeshi, Hokkaido, northern Japan. *Mycoscience* 46: 287e293.

White TJ, Bruns T, Lee SB, Taylor JW, 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky J, White T (eds), PCR protocols: a guide to methods and application. Academic Press, San Diego, pp 315e322.

Zhang Y, Crous PW, Schoch CL, Hyde KD, 2012. Pleosporales. *Fungal Diversity* 53: 1e221.

Zhang Y, Schoch CL, Fournier J, Crous PW, de Gruyter J, Woudenberg JHC, Hirayama K, Tanaka K, Pointing SB, Spatafora JW, Hyde KD, 2009a. Multi-locus phylogeny of Pleosporales: a taxonomic, ecological and evolutionary reevaluation. *Studies in Mycology* 64: 85e102.

Zhang Y, Wang HK, Fournier J, Crous PW, Jeewon R, Pointing SB, Hyde KD, 2009b. Towards a phylogenetic clarification of Lophiostoma/Massarina and morphologically similar genera in the Pleosporales. *Fungal Diversity* 38: 225e251.