


## The Face of Diabetes in Robeson County

What do we know about it?






- Diabetes affects 13.3% of our residents
- This is 4x greater than the state and 5x greater than the USA
- Lowry (1999) found <sup>1</sup>/<sub>2</sub> of everybody (Native Americans) in her study had diabetes
- Most of us are overweight = at risk



 Few residents use CAM (complementary alternative medicines) to manage diabetes • Few residents use diabetes specialists; most use primary care doctors.




## Female


 Diabetes mortality rate is highest among women living in eastern NC

## Male

 African American males are 2.3 times more likely to die from diabetes than European American men.



- Physical activity in rural Native and African American elders with diabetes is limited.
- Fifty percent (50%) of Lumbee women in a study were overweight or obese.
- Depression is twice as common among local residents with diabetes.



•

- About nine percent of Lumbee Indians over 18 years of age in Robeson County have been diagnosed with diabetes, compared to a statewide diabetes rate of about five percent, according to Ronny Bell, Ph.D., M.S., of Wake Forest University School of Medicine.
- "This higher rate of diabetes among the Lumbee validates what many people in this area believe -that diabetes is a major concern for Lumbees," Bell said. "We also know that in the general population, about one-third of people who actually have diabetes have not been diagnosed, so this rate actually misses quite a few people."

