

## REFERENCES

- ANDERSEN, R.A. (1982). A light and electron microscopical investigation of *Ochromonas sphaerocystis* Matvienko (Chrysophyceae): the statospore, vegetative cell and its peripheral vesicles. *Phycologia*, **21**: 390-398.
- ANDERSEN, R.A. (1987) Synurophyceae classis nov., a new class of algae. *Amer. J. Bot.*, **74**: 337-353.
- ANDERSEN, R.A. (1989). Absolute orientation of the flagellar apparatus of *Hibberdia magna* comb. nov. (Chrysophyceae). *Nord. J. Bot.*, **8**: 653-669.
- ANDERSEN, R.A. (1990). The three-dimensional structure of the flagellar apparatus of *Chrysosphaerella brevispina* (Chrysophyceae) as viewed by high voltage electron microscopy stereo pairs. *Phycologia*, **29**: 86-97.
- ANDERSEN, R.A., BARR, D.J.S., LYNN, D.H., MELKONIAN, M., MOESTRUP, Ø. & SLEIGH, M.A. (1991). Terminology and nomenclature of the cytoskeletal elements associated with the flagellar/ciliary apparatus in protists. *Protoplasma*, **164**: 1-8.
- ANDERSEN, R.A., SAUNDERS, G.W., PASKIND, M.P. & SEXTON, J.P. (1993). Ultrastructure and 18S rRNA gene sequence for *Pelagomonas calceolata* gen. et sp. nov. and the description of a new algal class, the Pelagophyceae classis nov. *J. Phycol.*, **29**: 701-715.
- ANDERSEN, R.A., MORTON, S.L. & SEXTON, J.P. (1997). CCMP - Provasoli-Guillard National Center for Culture of Marine Phytoplankton: List of Strains. *J. Phycol.* (suppl.), **33**: 1-75.
- ANDERSON, R.A., VAN de PEER, Y., POTTER, D., SEXTON, J.P., KAWACHI, M., & LaJEUNESSE, T. (1999). Phylogenetic analysis of the SSU rRNA from members of the Chrysophyceae. *Protist*, **150**: 71-84.
- ANDERSEN, R.A. & PREISIG, H.R. (2000). Class Synurophyceae. In *The Illustrated Guide to the Protozoa*, 2<sup>nd</sup> Edition (Lee, J.J., Leedale, G.F. & Bradbury, P., editors), 759-774. Allen Press Inc., Lawrence.
- ANDERSEN, R.A., POTTER, D. & BAILEY, J.C. (2002). *Pinguiococcus pyrenoidosus* gen. et. sp. nov. (Pinguiophyceae), a new marine coccoid alga. *Phycol. Res.*, **50**: 57-65.
- BAILEY, J.C., BIDIGARE, R.R., CHRISTENSEN, S.J., & ANDERSEN, R.A. (1998). Phaeothamniophyceae classis nova: A new lineage of chromophytes based upon photosynthetic pigments, *rbcL*, sequence analysis and ultrastructure. *Protist*, **149**: 245-263.

- BELCHER, J.H. & SWALE, E.M.F. (1972a). Some features of the microanatomy of *Chrysococcus cordiformis* Naumann. *Br. phycol. J.*, **7**: 53-59.
- BELCHER, J.H. & SWALE, E.M.F. (1972b). The morphology and fine structure of the colourless colonial flagellate *Anthophysa vegetans* (O.F. Müller) Stein. *Br. phycol. J.*, **7**: 335-346.
- BELCHER, J.H. (1974). *Chrysosphaera magna* sp. nov., a new coccoid member of the Chrysophyceae. *Br. phycol. J.*, **9**: 139-144.
- BELCHER, J.H. & SWALE, E.M.F. (1976). *Spumella elongata* (Stokes) comb. nov., a colourless flagellate from soil. *Arch. Protistenkd.*, Band **118**: 215-220.
- BJØRNLAND, T. & LIAAEN-JENSEN, S. (1989). Distribution patterns of carotenoids in relation to chromophyte phylogeny and systematics. In *The Chromophyte Algae: Problems and Perspectives* (Green, J.C., Leadbeater, B.S.C. & Diver, W.L., editors), 37-60. Clarendon Press, Oxford.
- BODDI, S., BIGAZZI, M. & SARTONI, G. (1999). Ultrastructure of vegetative and motile cells, and zoosporogenesis in *Chrysonephos lewissi* (Taylor) Taylor (Sarcinochrysidales, Pelagophyceae) in relation to taxonomy. *Eur. J. Phycol.* **34**: 297-306.
- BOUCK, G.B. (1969). Extracellular microtubules. The origin, structure, and attachment of flagellar hairs in *Fucus* and *Ascophyllum* anterozoids. *J. Cell Biol.*, **40**: 446-460.
- BOUCK, G.B. (1971). The structure, origin, isolation and composition of the tubular mastigonemes of the *Ochromonas* flagellum. *J. Cell Biol.*, **50**: 362-384.
- BOUCK, G.B. (1972). Architecture and assembly of mastigonemes. In *Advances in Cell and Molecular Biology*, Vol. 2 (DuPraw, D.J., editor), 237-271. Academic Press, New York.
- CAVALIER-SMITH, T., CHAO, E.E. & ALLSOPP, M.T.E.P. (1995). Ribosomal RNA evidence for chloroplast loss with Heterokonta: pedinellid relationships and a revised classification of ochristan algae. *Arch. Protistenkd.*, **145**: 209-220.
- CAVALIER-SMITH, T., CHAO, E.E., THOMPSON, C.E. & HOURIHANE, S.L. (1995/96). *Oikomonas*, a distinctive zooflagellate related to chrysomonads. *Arch. Protistenkd.*, **146**: 273-279.
- CHEN, L.L. & HAINES, T.H. (1975). The flagellar membrane of *Ochromonas danica*. *J. Biol. Chem.*, **251**: 1828-1834.
- CLARKE, K.J. & PENNICK, N.C. (1975). *Syncrypta glomerifera* sp. nov., a marine member of the Chrysophyceae bearing a new form of scale. *Br. phycol. J.*, **10**: 363-370.

- CLAYTON, M.N. (1989). Brown algae and chromophyte phylogeny. In *The Chromophyte Algae: Problems and Perspectives* (Green, J.C., Leadbeater, B.S.C. & Diver, W.L., editors), 229-253. Clarendon Press, Oxford.
- CLAYTON, M.N. (1990). Phylum Phaeophyta. In *Handbook of Protoctista* (Margulis, L., Corliss, J., Melkonian, M. & Chapman, D., editors), 698-714. Jones & Bartlett Publ., Boston.
- DAUGBJERG, N. (1996). *Mesopedinella arctica* gen. et sp. nov. (Pedinellales, Dictyochophyceae) I: fine structure of a new marine phytoflagellate from arctic Canada. *Phycologia*, **35**: 435-445.
- DAUGBJERG, N. & ANDERSEN, R.A. (1997a). A molecular phylogeny of the heterokont algae based on analyses of chloroplast-encoded *rbcL* sequence data. *J. Phycol.*, **33**:1031-1041.
- DAUGBJERG, N. & ANDERSEN, R.A. (1997b). Phylogenetic analyses of the *rbcL* sequences from haptophytes and heterokont algae suggest their chloroplasts are unrelated. *Mol. Biol. Evol.*, **14**: 1242-1251.
- DAUGBJERG, N. & GUILLOU, L. (2001). Phylogenetic analyses of the Bolidophyceae (Heterokontophyta) using *rbcL* gene sequences support their sister group relationship to diatoms. *Phycologia*, **40**:153-161.
- DOP, A.J. & VAN OERS, F. (1979). On the taxonomic position of *Phaeobotrys solitaria* Ettl (Chrysophyceae). *Acta Bot. Neerl.*, **28**: 459-466.
- DOYLE, J.J. & DOYLE, J.L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* **19**: 11-15.
- ENTWISLE, T.J. & ANDERSEN, R.A. (1990). A re-examination of *Tetrasporopsis* (Chrysophyceae) and the description of *Dermatochrysis* gen. nov. (Chrysophyceae): a monostromatic alga lacking cell walls. *Phycologia*, **29**: 263-274.
- ETTL, H. (1978). Süßwasserflora von Mitteleuropa. Band 3, Teil 1. Xanthophyceae. Gustav Fisher Verlag, Stuttgart.
- FOTT, B. (1956). Algenkunde. VEB Gustav Fischer Verlag, Jena.
- GRAHAM, L.E. & WILCOX, L.W. (2000). *Algae*. Prentice Hall, Upper Saddle River, NJ.
- GILBERT, D. (1994). SeqApp, v. 1.9a169, available via anonymous ftp at ftp.bio.indiana.edu

GUILLOU, L., CHRÉTIENNOT-DINET, M.-J., MEDLIN, L.K., CLAUSTRE, H., LOISEAUX-DE GOËR, S. & VAULOT, D. (1999). *Bolidomonas*: a new genus with two species belonging to a new algal class, the Bolidophyceae (Heterokonta). *J. Phycol.*, **35**: 368-381.

HEYWOOD, P. (1972). Structure and origin of flagellar hairs in *Vacuolaria virescens*. *J. Ultrastruct. Res.*, **39**: 608-623.

HEYWOOD, P. (1973). Intracisternal microtubules and flagellar hairs of *Gonyostomum semen* (Ehrenb.) Diesing. *Br. phycol. J.*, **8**: 43-46.

HEYWOOD, P. (1980). Chloromonads. In *Phytoflagellates* (Cox, E.R., ed.), 351-377. Elsevier North Holland, Inc.

HEYWOOD, P. & LEEDALE, G.F. (2000). Order Raphidomonadida. In *The Illustrated Guide to the Protozoa*, 2<sup>nd</sup> Edition (Lee, J.J., Leedale, G.F. & Bradbury, P., editors), 744-750. Allen Press Inc., Lawrence.

HIBBERD, D.J. (1970). Observations on the cytology and ultrastructure of *Ochromonas tuberulatus* sp. nov. (Chrysophyceae), with special reference to the discobolocysts. *Br. phycol. J.*, **5**: 119-143.

HIBBERD, D.J. (1971). Observations on the cytology and ultrastructure of *Chrysamoeba radians* Klebs (Chrysophyceae). *Br. phycol. J.*, **6**: 207-223.

HIBBERD, D.J. & LEEDALE, G.F. (1971). Cytology and ultrastructure of the Xanthophyceae. II. The zoospore and vegetative cell of coccoid forms, with special reference to *Ophiocytium majus*. *Br. phycol. J.*, **6**: 1-23.

HIBBERD, D.J. & LEEDALE, G.F. (1972). Observations on the cytology and ultrastructure of the new algal class, Eustigmatophyceae. *Ann. Bot.*, **36**: 49-71.

HIBBERD, D.J. (1976). The ultrastructure and taxonomy of the Chrysophyceae and Prymnesiophyceae (Haptophyceae): a survey with some new observations on the Chrysophyceae. *Bot. J. Linn. Soc.*, **72**: 55-80.

HIBBERD, D.J. (1977). The cytology and ultrastructure of *Chrysonebula holmesii* Lund (Chrysophyceae), with special reference to the flagellar apparatus. *Br. phycol. J.*, **12**: 369-383.

HIBBERD, D.J. (1980). Xanthophytes. In *Phytoflagellates* (Cox, E.R. editor). 243-271. Elsevier North-Holland, New York, Amsterdam, Oxford.

HIBBERD, D.J. (1990). Phylum Xanthophyta. In *Handbook of Protoctista* (Margulis, L., Corliss, J., Melkonian, M. & Chapman, D., editors), 686-697. Jones & Bartlett Publ., Boston.

- JEFFREY, S.W. (1989). Chlorophyll *c* pigments and their distribution in the chromophyte algae. In *The Chromophyte Algae: Problems and Perspectives* (Green, J.C., Leadbeater, B.S.C. & Diver, W.L., editors), 13-36. Clarendon Press, Oxford.
- KAPRAUN, D.F., DUTCHER, J.A. & LOPEZ-BAUTISTA, J. (1992). Nuclear genome characterization of the carrageenophyte *Agardhiella subulata* (Rhodophyta). *J. Appl. Phycol.* **4**: 129-137.
- KAWACHI, M., INOUE, I., HONDA, D., O'KELLY, C.J., BAILEY, J.C., BIDIGARE, R.R. & ANDERSEN, R.A. (2002). The Pinguiphyceae *classis nova*, a new class of photosynthetic stramenopiles whose members produce large amounts of omega-3 fatty acids. *Phycol. Res.*, **50**: 31-47.
- KAWAI, H., MAEBA, S., SASAKI, H. OKUDA, K. & HENRY, E.C. (2003). *Schizocladia ischiensis*: a new filamentous marine chromophyte belonging to a new class, Schizocladiphyceae. *Protist*, **154**: 211-228.
- KRISTIANSEN, J. (1990). Phylum Chrysophyta. In *Handbook of Protozoology* (Margulis, L., Corliss, J., Melkonian, M. & Chapman, D., editors), 438-453. Jones & Bartlett Publ., Boston.
- LEEDALE, G.F., LEADBEATER, B.S.C. & MASSALSKI, A. (1970). The intracellular origin of flagellar hairs in the Chrysophyceae and Xanthophyceae. *J. Cell Sci.*, **6**: 701-719.
- MIGNOT, J.P. (1976). Compléments à l'étude des chloromonades. Ultrastructure de *Chattonella subsalsa* Biecheler. Flagellé décaux saumâtre. *Protistologica*, **12**: 279-93.
- MOESTRUP, Ø. (1982). Flagellar structure in algae: a review, with new observations particularly on the Chrysophyceae, Phaeophyceae (Fucophyceae), Euglenophyceae and *Reckertia*. *Phycologia*, **21**: 427-528.
- MOESTRUP, Ø. & ANDERSEN, R.A. (1991). Organization of heterotrophic heterokonts. In *The Biology of Free-living Heterotrophic Flagellates* (Patterson, D.J. & Larsen, J., editors), 334-360. Clarendon Press, Oxford.
- MOESTRUP, Ø. (1995). Current status of the chrysophyte 'splinter groups': synurophytes, pedinellids, silicoflagellates. In *Chrysophyte Algae* (Sandgren, C.D., Smol, J.P. & Kristiansen, J., editors), 75-91. Cambridge University Press, Cambridge.
- MOESTRUP, Ø. & O'KELLY, C.J. (2000). Class Silicoflagellata. In *The Illustrated Guide to the Protozoa*, 2<sup>nd</sup> Edition (Lee, J.J., Leedale, G.F. & Bradbury, P., editors), 775-782. Allen Press Inc., Lawrence.
- MÜLLER, O.F. (1786). *Animalcula Infusoria Fluvialia et Marina*. Haniae, Copenhagen.

- O'KELLY, C.J. (1989). The evolutionary origin of the brown algae: information from studies of motile cell ultrastructure. In *The Chromophyte Algae: Problems and Perspectives* (Green, J.C., Leadbeater, B.S.C. & Diver, W.L., editors), 255-278. Oxford. of DNA substitution. *Bioinformatics*, **14**: 817-818.
- O'KELLY, C.J. & WUJEK, D.E. (2001). Cell structure and asexual reproduction in *Lagynion delicatulum* (Stylococcaceae, Chrysophyceae). *Eur. J. Phycol.*, **36**: 51-59.
- PÉTERFI, L.S. (1969). The fine structure of *Poterioochromonas malhamensis* (Pringsheim) comb. nov. with special reference to the lorica. *Nova Hedwigia*, **17**:93-103.
- PIPES, L.D., LEEDALE, G.F. & TYLER, P.A. (1981). Ultrastructure of *Tessellaria volvocina* (Synurophyceae). *Br. phycol. J.*, **26**: 259-278.
- POSADA, D. & CRANDALL, K.A. (1998). Modeltest: testing the model
- PREISIG, H.R. & HIBBERD, D.J. (1983). Ultrastructure and taxonomy of *Paraphysomonas* (Chrysophyceae) and related genera 3. *Nord. J. Bot.*, **3**: 695-723.
- PREISIG, H.R. & WILHELM, C. (1989). Ultrastructure, pigments and taxonomy of *Botryochloropsis similis* gen. et sp. nov. (Eustigmatophyceae). *Phycologia*, **28**: 61-69.
- PREISIG, H.R. (1995). A modern concept of chrysophyte classification. In *Chrysophyte Algae: Ecology, phylogeny and development* (Sandgren, C.D., Smol, J.P., & Kristiansen, J., editors), 46-74. Cambridge Univ Press, Cambridge.
- PREISIG, H.R. & ANDERSEN, R.A. (2000). Chrysomonada. In *The Illustrated Guide to the Protozoa*, 2<sup>nd</sup> Edition (Lee, J.J., Leedale, G.F. & Bradbury, P., editors), 693-730. Allen Press Inc., Lawrence.
- PRINGSHEIM, E.G. (1946). On iron flagellates. *Phil. Trans.*, **232**: 311-342.
- PRINGSHEIM, E.G. (1952). On the nutrition of *Ochromonas*. *J. Microscopical Sci.*, **93**: 71-96.
- REYNOLDS, E.S. (1963). The use of lead citrate at high pH as an electron opaque stain in electron microscopy. *J. Cell. Biol.*, **17**:208-212.
- RIETH, A. (1980). Xanthophyceae. In *Xanthophyceae Süßwasserflora von Mitteleuropa*. (Ettl, H., Gerlott, J., & Heynig, H. editors) 147. Gustav Fisher Verlag, Stuttgart.
- SANTOS, L.M.A. (1996). The Eustigmatophyceae: actual knowledge and research perspectives. *Nova Hedwigia*, **112**: 391-405.

SAUNDERS, G.W. & KRAFT, G.T. (1994). Small-subunit rRNA gene sequences from representatives of selected families of the Gigartinales and Rhodymeniales (Rhodophyta) Evidence for the Plocamiales *ord. nov.* *Can. J. Bot.* **72**: 1250-1263.

SCHNEPF, E., NIEMANN, A. & WILHELM, C. (1995/96). *Pseudostaurastrum limneticum*, a eustigmatophycean alga with astigmatic zoospores: morphogenesis, fine structure, pigment composition and taxonomy. *Arch. Protistenkd.*, **146**: 237-249.

SPURR, R.A. (1969). A low viscosity epoxy resin embedding medium for electron microscopy. *J. Ultrastruct. Res.*, **26**:31-34.

STARMACH, K (1985). Chrysophyceae und Haptophyceae. In Ettl H, Gerloff J, Heynig H, Mollenhauer D(eds) Süßwasserflora von Mitteleuropa. Gustav Fischer Verlag, Stuttgart, pp 1–515.

SWOFFORD, D.L. (2002) PAUP\*. Phylogenetic Analysis Using Parsimony (\*and Other Methods). Version 4. Sinauer Associates, Sunderland, Massachusetts.

THOMSEN, H.A. (1988). Ultrastructural studies of the flagellate and cyst stages of *Pseudopedinella tricostata* (Pedinellales, Chrysophyceae). *Br. phycol. J.*, **23**: 1-16.

TRENKWALDER, H. (1975). Neue Bodenalgen aus Föhrenwäldern im Raum von Brixen (Südtirol, Italien). *Ber. nat.-med. Ver. Innsbruck*, Band **62**: 7-19.

WUJEK, D.E. (1996). Chrysophyte ultrastructure and taxonomy: a mini review. In *Cytology, Genetics and Molecular Biology of Algae* (Chaudhary, B.R. & Agrawal, S.B., editors), 21-36. SPB Academic Publishing, Amsterdam, The Netherlands.