

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/320090100>

On the class of functions having infinite limit on a given set

Article in *Colloquium Mathematicum* · January 1994

DOI: 10.4064/cm-67-2-177-180

CITATION

1

READS

2

2 authors:



János T. Tóth

60 PUBLICATIONS 136 CITATIONS

SEE PROFILE



Laszlo Zsilinszky

University of North Carolina at Pembroke

42 PUBLICATIONS 152 CITATIONS

SEE PROFILE

ON THE CLASS OF FUNCTIONS HAVING
INFINITE LIMIT ON A GIVEN SET

BY

J. TÓTH AND L. ZSILINSZKY (NITRA)

Introduction. Given a topological space X and a real function f on X define

$$L_f(X) = \{x \in X : \lim_{t \rightarrow x} f(t) = +\infty\}.$$

According to [1] for a linear set A there exists a function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $A = L_f(\mathbb{R})$ if and only if A is a countable G_δ -set. Our purpose is to prove a similar result in a more general setting and to investigate the cardinality and topological properties of the class of functions $f : X \rightarrow \mathbb{R}$ for which $L_f(X)$ equals a given non-empty, countable G_δ -set.

We will need some auxiliary notions and notations. Denote by \bar{E} and E^c , respectively, the closure and the set of all condensation points of a subset E of a topological space, and by $\text{card } E$ its cardinality. Denote by \mathcal{F} the space \mathbb{R}^X .

A topological space X is called a *Fréchet space* if for every $E \subset X$ and every $x \in \bar{E}$ there exists a sequence in E converging to x (cf. [2]). Every first-countable space is a Fréchet space ([2], p. 78), but there exists a Fréchet space that is not first-countable ([2], p. 79).

A topological space X is said to be *hereditarily Lindelöf* if for each $E \subset X$ every open cover of E has a countable refinement. A well-known property of these spaces is as follows ([4], p. 57):

LEMMA 1. *If X is a hereditarily Lindelöf space, then $E \setminus E^c$ is countable for each $E \subset X$.*

Main results. Using Lemma 1 it can be shown similarly to [1] that for a Hausdorff, hereditarily Lindelöf space X having no isolated points, $L_f(X)$ is a countable G_δ -set for every $f \in \mathcal{F}$. We will be interested in the reverse problem, namely to find, for every non-empty, countable G_δ -set $A \subset X$, a function $f \in \mathcal{F}$ for which $L_f(X) = A$.

1991 *Mathematics Subject Classification*: Primary 54C30.

In what follows X will be a Fréchet, Hausdorff, hereditarily Lindelöf space such that $X = X^c$. Let A be a given non-empty, countable G_δ -subset of X . Define

$$\mathcal{S} = \{f \in \mathcal{F} : L_f(X) = A\}.$$

THEOREM 1. *The set \mathcal{S} is non-empty.*

Proof. Let $A = \{a_1, a_2, \dots\} \subset X$, $A = \bigcap_{n=1}^{\infty} G_n$ where $G_1 = X$, G_n is open in X , $G_{n+1} \subsetneq G_n$ ($n \in \mathbb{N}$). We can assume that $F_n = G_n \setminus G_{n+1}$ is uncountable for each $n \in \mathbb{N}$. Put $H = \bigcup_{n=1}^{\infty} (F_n \cap F_n^c)$.

According to Lemma 1 the set $B = (\bigcup_{n=1}^{\infty} F_n) \setminus H$ is countable, since $B \subset \bigcup_{n=1}^{\infty} (F_n \setminus F_n^c)$. Write $B = \{b_1, b_2, \dots\}$. Observe that $A \cap \bigcup_{n=1}^{\infty} F_n = \emptyset$, so

$$X \setminus H = \left(A \cup \bigcup_{n=1}^{\infty} F_n \right) \setminus H = A \cup B.$$

Therefore $B \subset \overline{H}$ (since $B \subset X^c$), hence for each $k \in \mathbb{N}$ there exists a sequence $c_i^{(k)} \in H$ ($i \in \mathbb{N}$) converging to b_k . Set $C = \bigcup_{i,k \in \mathbb{N}} \{c_i^{(k)}\}$. Define a function $f \in \mathcal{F}$ as follows:

$$\begin{aligned} f(a_k) &= k && \text{for all } k \in \mathbb{N}, \\ f(c_i^{(k)}) &= k && \text{for all } i, k \in \mathbb{N}, \\ f(x) &= n && \text{for all } x \in F_n \setminus C, n \in \mathbb{N}. \end{aligned}$$

We will prove that $L_f(X) = A$.

First choose $x \in X \setminus A$. Then either $x \in B$ or $x \in H$. If $x \in B$ then $x = b_k$ for some $k \in \mathbb{N}$, and consequently $x \notin L_f(X)$ since $\lim_{i \rightarrow \infty} f(c_i^{(k)}) = k$. If $x \in H$ then $x \in F_m^c$ for some $m \in \mathbb{N}$, so there exists a directed set Σ and a net $\{x_\sigma : \sigma \in \Sigma\}$ in $F_m \setminus C$ converging to x . Thus again $x \notin L_f(X)$ since $\lim_\sigma f(x_\sigma) = m$.

Finally, suppose $x \in A$. Take an arbitrary $n \in \mathbb{N}$. Then $x \in G_n$. The space X is Hausdorff, so there is a neighbourhood S_1 of x which contains no member of the sequence $\{c_i^{(k)}\}_{i=1}^{\infty}$ for all $1 \leq k \leq n$ (notice that $c_i^{(k)} \rightarrow b_k \notin A$ as $i \rightarrow \infty$). Further, there exists a neighbourhood S_2 of x containing none of a_1, \dots, a_n except possibly x . It is now not hard to see that $f(t) \geq n$ for each $t \in G_n \cap S_1 \cap S_2$, $t \neq x$, whence $x \in L_f(X)$. ■

Remark 1. If X is a Hausdorff, second-countable, Baire space with no isolated points (in particular, if X is a separable, complete metric space with no isolated points) then Theorem 1 holds. Indeed, in this case every non-empty open subset of X is uncountable (see [3], Proposition 1.29) and thus $X = X^c$; further, second-countable spaces are Fréchet and hereditarily Lindelöf.

THEOREM 2. *We have $\text{card } \mathcal{S} = \text{card}(\mathcal{F} \setminus \mathcal{S}) = 2^{\text{card } X}$.*

Proof. Let $f \in \mathcal{S}$ (see Theorem 1). Using the notation of Theorem 1 put $\alpha_n = \text{card}(X \setminus G_n)$ ($n \in \mathbb{N}$) and $\alpha = \text{card}(X \setminus A) = \text{card } X$ (X is uncountable). Then $\{\alpha_n\}_{n=1}^{\infty}$ is a non-decreasing sequence of infinite ordinals converging to α (in the order topology; see [4]). Fix $n \in \mathbb{N}$. For every $M \subset X \setminus G_n$ define the function $f_M = \max\{1, f\} \cdot \chi_{X \setminus M}$, where $\chi_{X \setminus M}$ is the characteristic function of $X \setminus M$.

It is not hard to see that $f_M \neq f_N$ and $f_M, f_N \in \mathcal{S}$ for any different subsets M, N of the closed set $X \setminus G_n$. Thus $\text{card } \mathcal{S} \geq 2^{\alpha_n}$. Since α is a limit ordinal we have $\text{card } \mathcal{S} \geq \sup\{2^{\alpha_n} : n \in \mathbb{N}\} = 2^{\alpha} = 2^{\text{card } X}$. On the other hand, making allowance for the uncountability of X we get $\text{card } \mathcal{S} \leq \text{card } \mathcal{F} = (\text{card } \mathbb{R})^{\text{card } X} = 2^{\text{card } X}$.

To show that $\text{card}(\mathcal{F} \setminus \mathcal{S}) = 2^{\text{card } X}$ it suffices to notice that $\chi_B \in \mathcal{F} \setminus \mathcal{S}$ for any $B \subset X$. Hence $2^{\text{card } X} \leq \text{card}(\mathcal{F} \setminus \mathcal{S}) \leq \text{card } \mathcal{F} = 2^{\text{card } X}$. ■

To be able to investigate \mathcal{S} from the topological point of view introduce the sup-metric d on \mathcal{F} :

$$d(f, g) = \min\{1, \sup_{x \in \mathbb{R}} |f(x) - g(x)|\}, \quad \text{where } f, g \in \mathcal{F}.$$

It is known that (\mathcal{F}, d) is a complete metric space.

THEOREM 3. *The class \mathcal{S} is simultaneously open and closed in \mathcal{F} .*

Proof. If $f, g \in \mathcal{F}$ and $d(f, g) < 1$ then $L_f(X) = L_g(X)$. So if $f \in \mathcal{S}$ (resp. $f \in \mathcal{F} \setminus \mathcal{S}$) then the open 1-ball around f is in \mathcal{S} (resp. in $\mathcal{F} \setminus \mathcal{S}$). ■

THEOREM 4. *Both \mathcal{S} and $\mathcal{F} \setminus \mathcal{S}$ are of second category in \mathcal{F} .*

Proof. According to Theorems 2 and 3, \mathcal{S} and $\mathcal{F} \setminus \mathcal{S}$ are non-empty open sets, and consequently they are of second category in the complete metric space (\mathcal{F}, d) . ■

Remark 2. In the light of Theorems 2 and 4 it is worth noticing that neither \mathcal{S} nor $\mathcal{F} \setminus \mathcal{S}$ is dense in \mathcal{F} . Actually, if, say, \mathcal{S} were dense in \mathcal{F} then in view of Theorem 3 it would be a residual set in \mathcal{F} and hence $\mathcal{F} \setminus \mathcal{S}$ of first category in \mathcal{F} .

THEOREM 5. *We have $\mathcal{S} \subset \mathcal{S}^c$ and $\mathcal{F} \setminus \mathcal{S} \subset (\mathcal{F} \setminus \mathcal{S})^c$.*

Proof. Let $f \in \mathcal{F}$ and $0 < \varepsilon < 1$. For $0 < \eta < \varepsilon$ define $f_{\eta}(x) = f(x) + \eta$ ($x \in X$). Then $d(f, f_{\eta}) = \eta < \varepsilon$ for all $\eta \in (0, \varepsilon)$; furthermore, $f_{\eta} \in \mathcal{S}$ if and only if $f \in \mathcal{S}$ ($0 < \eta < \varepsilon$). ■

Remark 3. It is easy to see that the set

$$\mathcal{S}' = \{f \in \mathcal{F} : \lim_{t \rightarrow x} f(t) = -\infty \text{ if and only if } x \in A\}$$

also has the properties established in Theorems 1–5 for \mathcal{S} .

REFERENCES

- [1] R. Bumcrot and M. Sheingorn, *Variations on continuity: sets of infinite limit*, Math. Mag. 47 (1974), 41-43.
- [2] R. Engelking, *General Topology*, PWN, Warszawa, 1977.
- [3] R. C. Haworth and R. A. McCoy, *Baire spaces*, Dissertationes Math. 141 (1977).
- [4] J. L. Kelley, *General Topology*, Springer, New York, 1975.

DEPARTMENT OF MATHEMATICS
COLLEGE OF EDUCATION
FARSKÁ 19
94974 NITRA, SLOVAKIA
E-mail: ZSILI@UNITRA.SK

Reçu par la Rédaction le 5.7.1993