

## SUPERPOROSITY IN A CLASS OF NON-NORMABLE SPACES

### INTRODUCTION

The concept of porous set was introduced by Dolženko in [D]. Since then it has been thoroughly investigated and diversely generalized (see [Za1] or [Re] for a survey). It is possible to define several notions concerning porosity also in metric spaces (see [Za1],[Re]). It is known that in Banach spaces the ideal of meager sets is strictly wider than that of the  $\sigma$ -porous sets ([Za1]). It is true also in closed non-locally compact convex subsets of a separable Banach space ([AB]). Recently it has been established in dense in itself completely metrizable spaces as well (cf. [Za3]).

The primary goal of the research presented in this paper is in the line of the above results, i.e. to compare  $\sigma$ -porous and meager sets, respectively in some non-normable spaces. Such an attempt was made in [TZs] where the space  $s$  of all real sequences endowed with the Fréchet metric

$$\rho_F(\{a_n\}_n, \{b_n\}_n) = \sum_n 2^{-n} \frac{|a_n - b_n|}{1 + |a_n - b_n|} \text{ where } \{a_n\}_n, \{b_n\}_n \in s$$

was scrutinized in this respect. This space is non-normable ([KG], Exercise 276) and it was shown in [TZs] e.g. that the set  $\{\{a_n\}_n \in s; \sum_n \Phi(a_n) \text{ converges}\}$  is  $\sigma$ -superporous in  $s$  for a residual family of functions  $\Phi$  in the space of all real functions furnished with the uniform topology.

It is the purpose of this paper to carry on these investigations generalizing results of [TZs] for the space  $\mathcal{M}$  of all measurable functions on an infinite  $\sigma$ -finite measure space  $(X, \mathcal{S}, \mu)$  endowed with the (metrizable) topology of convergence in measure on sets of finite measure (see [G]). We will show that results quite analogous to those of exposed in [TZs] for  $s$  hold in this generality as well. For instance, the set  $A(\Phi) = \{f \in \mathcal{M}; \int_X |\Phi \circ f| d\mu^* < +\infty\}$  is  $\sigma$ -superporous in  $\mathcal{M}$  for a broad class of functions  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$ , where  $\mu^*$  is the outer measure induced by  $\mu$  and  $\int_X^* h d\mu^*$  stands for the  $\mu^*$ -upper integral of the function  $h : X \rightarrow \mathbb{R}$  (see [F], Section 2.4).

Further we show that  $A(\chi_{\mathbb{R} \setminus M})$  is  $\sigma$ -superporous in  $\mathcal{M}$  for every  $\sigma$ -very porous set  $M \subset \mathbb{R}$  ( $\chi_{\mathbb{R} \setminus M}$  is the characteristic function of  $\mathbb{R} \setminus M$ ) and that  $A(\chi_{\mathbb{R} \setminus M})$  is meager in  $\mathcal{M}$  if  $M$  is meager at some point of  $\mathbb{R}$ . In particular,  $A(\chi_{\mathbb{R} \setminus M})$  is meager in  $(s, \rho_F)$  if and only if  $M$  is meager at some point of  $\mathbb{R}$ .

This could provide a method for relating meager non- $\sigma$ -porous subsets of  $\mathbb{R}$  to meager non- $\sigma$ -porous subsets of  $\mathcal{M}$  (resp.  $s$ ) if the porosity of  $A(\chi_{\mathbb{R} \setminus M})$  in  $\mathcal{M}$  (resp.  $s$ ) could be characterized in terms of  $M \subset \mathbb{R}$ .

It is worth noticing here that a more familiar metrization of  $\mathcal{M}$  by the metric

$$m(f, g) = \inf\{\varepsilon > 0; \mu(\{x \in X; |f(x) - g(x)| \geq \varepsilon\}) < \varepsilon\} \quad (f, g \in \mathcal{M})$$

which coincides with the topology of convergence in measure on  $X$  (cf. [F], Section 2.3.8), yields a setting where our considerations are not feasible even for continuous  $\Phi$ 's. This question was studied in [Zs1].

### PRELIMINARIES

In the sequel  $(X, S, \mu)$  will be an infinite  $\sigma$ -finite measure space and  $\mu^*$  the outer measure induced by  $\mu$ . Without loss of generality we may suppose that  $X = \bigcup_{n=1}^{\infty} X_n$ , where  $\{X_n\}_{n=1}^{\infty}$  is a sequence of pairwise disjoint,  $S$ -measurable sets such that  $2 < \mu(X_n) < +\infty$  for each  $n \in \mathbb{N}$ .

Denote by  $\mathcal{M}$  (resp.  $\mathcal{M}_n$ ) the set of all  $S$ -measurable functions that are finite almost everywhere (abbr. a.e.) on  $X$  (on  $X_n$ ). We will identify members of  $\mathcal{M}$  provided they equal a.e. on  $X$ . If the sequence  $f_k \in \mathcal{M}$  ( $k \in \mathbb{N}$ ) converges in measure to  $f \in \mathcal{M}$ , write  $f_k \xrightarrow{\mu} f$ .

Denote by  $\mathcal{F}_m$  the space of all functions  $\Phi : \mathbb{R} \rightarrow \mathbb{R}$  such that  $\Phi \circ f \in \mathcal{M}$  for all  $f \in \mathcal{M}$ . It is known that  $\mathcal{F}_m$  contains the class of Borel-measurable functions. Observe that  $\mathcal{F}_m$  is a closed subspace of the complete metric space  $(\mathcal{F}, d)$ , where  $\mathcal{F} = \mathbb{R}^{\mathbb{R}}$  and

$$d(\Phi, \Psi) = \min\{1, \sup_{t \in \mathbb{R}} |\Phi(t) - \Psi(t)|\} \quad (\Phi, \Psi \in \mathcal{F}).$$

Indeed, if a sequence  $\Phi_n \in \mathcal{F}_m$  ( $n \in \mathbb{N}$ )  $d$ -converges to  $\Phi \in \mathcal{F}$  then  $\Phi_n \circ f \in \mathcal{M}$  converges pointwise to  $\Phi \circ f$  (for all  $f \in \mathcal{M}$ ), thus  $\Phi \circ f \in \mathcal{M}$  and consequently  $\Phi \in \mathcal{F}_m$ . It follows that  $(\mathcal{F}_m, d)$  is a complete metric space.

For  $\Phi \in \mathcal{F}$  and  $p \in \mathbb{N}$  define

$$A(\Phi) = \{f \in \mathcal{M}; \int_X^* |\Phi \circ f| d\mu^* < +\infty\} \text{ and}$$

$$A_p(\Phi) = \{f \in \mathcal{M}; \int_X^* |\Phi \circ f| d\mu^* \leq p\},$$

where  $\int_X^* f d\mu^*$  is the upper integral of  $f$  with respect to  $\mu^*$  (see [F], Section 2.4).

For  $f, g \in \mathcal{M}$  and  $n \in \mathbb{N}$  define

$$\rho_n(f, g) = \int_{X_n} \frac{|f - g|}{1 + |f - g|} d\mu$$

$$\rho(f, g) = \sum_{n=1}^{\infty} \frac{1}{2^n \mu(X_n)} \rho_n(f, g).$$

For  $i, j \in \mathbb{N}$  and  $M \subset \mathbb{R}$  denote

$$\tilde{A}_{i,j}(M) = \{f \in \mathcal{M}; \mu^*(f^{-1}(M) \cap X_i) \geq \frac{\mu(X_i)}{j}\},$$

$$A_{i,j}(M) = \{f|_{X_i}; f \in \tilde{A}_{i,j}(M)\} \text{ and } A_{i,0}(M) = \{f \in \mathcal{M}_i; \mu^*(f^{-1}(M)) = \mu(X_i)\}.$$

It is not hard to see that  $\rho$  (resp.  $\rho_n$ ) is a metric on  $\mathcal{M}$  (resp.  $\mathcal{M}_n$ ). It can be shown similarly as for  $(s, \rho_F)$  that  $(\mathcal{M}, \rho)$  is non-normable (see [Zs2]).

Convergence in measure implies  $\rho$ -convergence and the converse holds if and only if the underlying measure space is finite. More precisely we have:

**Lemma 1.** *Let  $f_k, f \in \mathcal{M}$  ( $k \in \mathbb{N}$ ). The following are equivalent:*

- (i)  $f_k \xrightarrow{\rho} f$ ;
- (ii)  $f_k \xrightarrow{\mu} f$  on every  $S$ -measurable set of finite measure;
- (iii)  $f_k|_{X_n} \xrightarrow{\rho_n} f|_{X_n}$  for all  $n \in \mathbb{N}$ .

*Proof.* For (i) $\Leftrightarrow$ (ii) see [G], Theorem 3. The equivalence (i) $\Leftrightarrow$ (iii) follows easily from [K] (Theorem 14, p.122).  $\square$

*Remark 1.* Observe that  $(\mathcal{M}_n, \rho_n)$  is a complete metric space for each  $n \in \mathbb{N}$  and the  $\rho_n$ -convergence of sequences from  $\mathcal{M}_n$  coincides with their convergence in measure on  $X_n$  ([Ha], Problem 42(4)). Further the equivalence (i) $\Leftrightarrow$ (iii) in the previous lemma actually yields that  $(\mathcal{M}, \rho)$  and the Tychonoff product  $\Pi_n(\mathcal{M}_n, \rho_n)$  are homeomorphic.

**Lemma 2.** (cf.[G])  *$(\mathcal{M}, \rho)$  is a complete metric space.*

Denote by  $B_d(y, r)$  the open ball about  $y \in Y$  with radius  $r > 0$  in the metric space  $(Y, d)$ . By  $B(x, r)$  we will denote the interval  $(x - r, x + r)$ , where  $x \in \mathbb{R}$ . For  $E \subset Y, y \in Y$  and  $r > 0$  define

$$\gamma(y, r, E) = \sup\{r' > 0; \exists y' \in Y \ B_d(y', r') \subset B_d(y, r) \setminus E\}.$$

We say that  $E$  is *porous* (*very porous*) at  $y$  if

$$\limsup_{r \rightarrow 0^+} \frac{\gamma(y, r, E)}{r} > 0 \quad (\liminf_{r \rightarrow 0^+} \frac{\gamma(y, r, E)}{r} > 0).$$

Further  $E$  is said to be *superporous* at  $y \in Y$  (see [Za1],[Za2]), if  $E \cup F$  is porous at  $y$  whenever  $F \subset Y$  is porous at  $y$ .

A set  $E \subset Y$  is said to be *globally very porous* if there exist constants  $0 < a_E < 1$  and  $r_0 > 0$  such that  $\gamma(y, r, E) > a_E r$  for every  $y \in E$  and  $0 < r < r_0$  ([Za1]).

We say that  $E$  is *superporous* (*very porous*) if it is superporous (*very porous*) at each of its points, further  $E$  is  $\sigma$ -superporous ( $\sigma$ -very porous) if it is a countable union of superporous (*very porous*) sets. Superporosity was defined in [Za2] in connection with the  $\mathcal{I}$ -density topology of Wilczynski and others (cf.[W]).

Note that superporosity implies very porosity as observed in [Za2] (see [Re], Corollary 8.15 as well) and  $\sigma$ -superporosity is equivalent to  $\sigma$ -very porosity which is further equivalent to  $\sigma$ -globally very porosity ([Re], Corollary 8.17).

We will denote by  $card Y$  and  $\mathcal{P}(Y)$  the cardinality and the power set, respectively of the set  $Y$ , further  $c$  will stand for the power of the continuum. Denote by  $|I|$  the length of the interval  $I \subset \mathbb{R}$ .

## MAIN RESULTS

**Lemma 3.** *Let  $\{I_q; q \in \mathbb{N}\}$  be an enumeration of open intervals with rational endpoints. Let  $\Phi_{pq} = p\chi_{I_q}$  for  $p, q \in \mathbb{N}$ . Then  $A_p(\Phi_{pq})$  is superporous in  $(\mathcal{M}, \rho)$  for every  $p, q \in \mathbb{N}$ .*

*Proof.* Choose  $p, q \in \mathbb{N}$  and denote by  $t_q$  the midpoint of  $I_q$ . Let  $f \in A_p(\Phi_{pq})$ . Suppose that  $F \subset \mathcal{M}$  is an arbitrary set porous at  $f$ . Then there exist sequences

$r_n, r'_n > 0$  ( $n \in \mathbb{N}$ ) and  $\alpha > 0$  such that  $\alpha r_n < r'_n < r_n < 2^{-n}$ , further we get an  $f_n \in \mathcal{M}$  such that

$$(1) \quad B_\rho(f_n, r'_n) \subset B_\rho(f, r_n) \setminus F.$$

Define  $p_n = \min\{k \in \mathbb{N}; 2^{-k} < r'_n\} + 1$  and  $\varepsilon_n = 2^{-p_n+1}$  for all  $n \in \mathbb{N}$ . Then we have

$$(2) \quad r'_n > \varepsilon_n \geq \frac{r'_n}{2}.$$

Denote  $E_{n1} = X_{p_n} \cap f_0^{-1}((t_q - \frac{1}{8}|I_q|, t_q + \frac{1}{8}|I_q|))$  and  $E_{n2} = X_{p_n} \setminus E_{n1}$  and define  $g_n = f_n \chi_{X \setminus X_{p_n}} + t_q \chi_{E_{n2}} + (t_q + \frac{1}{4}|I_q|) \chi_{E_{n1}} \in \mathcal{M}$ . It is clear that

$$(3) \quad |f_n(x) - g_n(x)| \geq \frac{1}{8}|I_q| \text{ for all } x \in X_{p_n}.$$

Since  $\rho(f_n, g_n) = \frac{1}{2^{p_n} \mu(X_{p_n})} \int_{X_{p_n}} \frac{|f_n - g_n|}{1 + |f_n - g_n|} d\mu$  then by the definition of  $\varepsilon_n, X_{p_n}$  and (3), respectively we get

$$(4) \quad \rho(f_n, g_n) < \frac{\varepsilon_n}{2},$$

$$(4') \quad \rho(f_n, g_n) > \frac{|I_q|}{8 + |I_q|} \cdot \frac{\varepsilon_n}{2}.$$

Put  $\delta_n = \frac{|I_q|}{16 + 2|I_q|} \rho(f_n, g_n)$  and pick an arbitrary  $h_n \in B_\rho(g_n, \delta_n)$ . Define

$$D_n = \{x \in X_{p_n}; |h_n(x) - g_n(x)| < \frac{4\delta_n}{\varepsilon_n - 4\delta_n}\} \text{ and } D_{n0} = X_{p_n} \setminus D_n.$$

Observe that  $D_n$  is well-defined, since by (4)  $\delta_n = \frac{|I_q|}{8 + |I_q|} \cdot \frac{\rho(f_n, g_n)}{2} < \frac{\varepsilon_n}{4}$ .

Then we have

$$\begin{aligned} \delta_n > \rho(h_n, g_n) &\geq \frac{1}{2^{p_n} \mu(X_{p_n})} \int_{D_{n0}} \frac{|h_n - g_n|}{1 + |h_n - g_n|} d\mu \geq \\ &\geq \frac{\varepsilon_n}{2\mu(X_{p_n})} \int_{D_{n0}} \frac{4\delta_n}{\varepsilon_n} d\mu = \frac{2\delta_n \mu(D_{n0})}{\mu(X_{p_n})}, \end{aligned}$$

thus  $\mu(D_{n0}) < \frac{1}{2}\mu(X_{p_n})$  hence  $\mu(D_n) \geq \frac{1}{2}\mu(X_{p_n}) > 1$ .

In view of (4) we get  $|h_n(x) - g_n(x)| < \frac{4\delta_n}{\varepsilon_n - 4\delta_n} < \frac{1}{8}|I_q|$  for every  $x \in D_n$ , so  $h_n(D_n) \subset (t_q - \frac{3}{8}|I_q|, t_q + \frac{3}{8}|I_q|)$  (see the definition of  $g_n$ ). Then  $\int_X^* |\Phi_{pq} \circ h_n| d\mu^* \geq \int_{D_n}^* |\Phi_{pq} \circ h_n| d\mu^* \geq p\mu(D_n) > p$ , so

$$(5) \quad h_n \in \mathcal{M} \setminus A_p(\Phi_{pq}).$$

Using (4) we get  $\varepsilon_n - \rho(f_n, g_n) > \frac{\varepsilon_n}{2} > \frac{\varepsilon_n}{2} \cdot \frac{|I_q|}{8 + |I_q|} > \delta_n$ , therefore  $B_\rho(g_n, \delta_n) \subset B_\rho(f_n, \varepsilon_n) \subset B_\rho(f_n, r'_n)$ . Then in virtue of (5) and (1) there holds

$$B_\rho(g_n, \delta_n) \subset B_\rho(f_n, r'_n) \setminus A_p(\Phi_{pq}) \subset B_\rho(f, r_n) \setminus (F \cup A_p(\Phi_{pq})).$$

From (4') and (2) we get

$$\gamma(f, r_n, F \cup A_p(\Phi_{pq})) \geq \delta_n \geq \left(\frac{|I_q|}{8 + |I_q|}\right)^2 \frac{\varepsilon_n}{2} \geq \left(\frac{|I_q|}{8 + |I_q|}\right)^2 \frac{r'_n}{4} > \left(\frac{|I_q|}{8 + |I_q|}\right)^2 \frac{\alpha}{4} r_n,$$

thus  $\limsup_{r \rightarrow 0^+} \frac{\gamma(f, r, F \cup A_p(\Phi_{pq}))}{r} \geq \left(\frac{|I_q|}{8 + |I_q|}\right)^2 \frac{\alpha}{4} > 0$ , which proves the porosity of  $F \cup A_p(\Phi_{pq})$  at  $f$ .  $\square$

**Theorem 1.** *Let  $\Phi \in \mathcal{F}$  be a function for which there exists  $t_0 \in \mathbb{R} \cup \{\pm\infty\}$  such that*

$$(6) \quad \liminf_{t \rightarrow t_0} |\Phi(t)| > 0.$$

*Then  $A(\Phi)$  is  $\sigma$ -superporous in  $(\mathcal{M}, \rho)$ .*

*Proof.* In view of (6) there exists  $\beta > 0$  and a bounded open interval  $I$  such that

$$(7) \quad |\Phi(t)| \geq \beta \text{ for all } t \in I.$$

Let  $\{J_k; k \in \mathbb{N}\}$  be a partition of  $I$  consisting of open intervals. Choose an  $f \in A(\Phi)$ . Then by (7) we have

$$\beta \sum_{k \in \mathbb{N}} \mu(f^{-1}(J_k)) = \beta \mu(f^{-1}(I)) \leq \int_X |\Phi \circ f| d\mu^* < p$$

for some  $p \in \mathbb{N}$ . Thus  $\mu(f^{-1}(J_k)) \leq 1$  for some  $k \in \mathbb{N}$  and hence  $\mu(f^{-1}(I_q)) \leq 1$  for some open interval  $I_q \subset J_k$  with rational endpoints. Consequently,

$$\int_X |\Phi_{pq} \circ f| d\mu^* = p \mu(f^{-1}(I_q)) \leq p,$$

so  $f \in A_p(\Phi_{pq})$ , whence  $A(\Phi) \subset \bigcup_{p,q \in \mathbb{N}} A_p(\Phi_{pq})$ , which concludes the proof by Lemma 3.

□

As the following results show, there are also functions  $\Phi$ , not necessarily satisfying (6), for which  $A(\Phi)$  is still  $\sigma$ -superporous (cf. Theorem 2):

**Lemma 4.** *Let  $M \subset \mathbb{R}$  be a globally very porous set. Then  $\tilde{A}_{i,j}(M)$  is superporous in  $(\mathcal{M}, \rho)$  for each  $i, j \in \mathbb{N}$ .*

*Proof.* According to the assumption on  $M$  there exist  $0 < a_M < 1$  and  $r_0 > 0$  such that

$$(8) \quad \gamma(x, r, M) > a_M r \text{ for all } x \in M \cup (\mathbb{R} \setminus \overline{M}) \text{ and all } 0 < r < r_0.$$

Choose  $f \in \tilde{A}_{i,j}(M)$  and a set  $F \subset \mathcal{M}$  which is porous at  $f$ . Then there exist  $\alpha > 0$ , sequences  $r_n, r'_n > 0$  and  $f_n \in \mathcal{M}$  such that  $r_n \searrow 0, \alpha r_n < r'_n < r_n < 2^{-i+1} \cdot \frac{3r_0}{1+r_0}$  and

$$(9) \quad B(f_n, r'_n) \subset B(f, r_n) \setminus F.$$

It is not hard to find  $b_{nk} \in \mathbb{R}$  ( $1 \leq k \leq m_n$ , where  $m_n \in \mathbb{N}$ ) and a partition  $\{D_{nk}; 1 \leq k \leq m_n\}$  of  $X_i$  such that for  $g_{n0} = f_n \chi_{X \setminus X_i} + \sum_{k=1}^{m_n} b_{nk} \chi_{D_{nk}} \in \mathcal{M}$  there holds

$$(10) \quad \rho(f_n, g_{n0}) < \frac{r'_n}{4}.$$

We can actually choose  $b_{nk} \in M \cup (\mathbb{R} \setminus \overline{M})$  for every  $1 \leq k \leq m_n$ .

Put  $\eta_n = \frac{2^i r'_n}{6 - 2^i r'_n}$ . Then  $\eta_n < r_0$ , so it follows from (8) that for each  $1 \leq k \leq m_n$  there exists  $b'_{nk} \in \mathbb{R}$  and  $r_{nk} > 0$  such that

$$(11) \quad a_M \eta_n \leq r_{nk} < \eta_n \text{ and } B(b'_{nk}, r_{nk}) \subset B(b_{nk}, \eta_n) \setminus M.$$

Define  $g_n = g_{n0} \chi_{X \setminus X_i} + \sum_{k=1}^{m_n} b'_{nk} \chi_{D_{nk}} \in \mathcal{M}$ . Then by (11) we have

$$\begin{aligned} \rho(g_{n0}, g_n) &\leq \frac{1}{2^i \mu(X_i)} \sum_{k=1}^{m_n} \left( \int_{D_{nk}} |b_{nk} - b'_{nk}| d\mu \right) = \frac{1}{2^i \mu(X_i)} \sum_{k=1}^{m_n} |b_{nk} - b'_{nk}| \mu(D_{nk}) \leq \\ &\leq \frac{1}{2^i \mu(X_i)} \eta_n \sum_{k=1}^{m_n} \mu(D_{nk}) = \frac{r'_n}{6 - 2^i r'_n} \leq \frac{r'_n}{4}, \end{aligned}$$

thus in view of (10)

$$(12) \quad \rho(f_n, g_n) \leq \rho(f_n, g_{n0}) + \rho(g_{n0}, g_n) \leq \frac{r'_n}{2}.$$

We have  $0 < a_M < 1 < 3j$ , thus  $\frac{r'_n}{2} > \frac{a_M r'_n}{6j}$ . Then putting  $\delta_n = \frac{a_M r'_n}{6j}$  we get by (12) that  $r'_n - \rho(f_n, g_n) \geq \frac{r'_n}{2} > \delta_n$ , so

$$(13) \quad B_\rho(g_n, \delta_n) \subset B_\rho(f_n, r'_n).$$

Choose  $h \in \tilde{A}_{i,j}(M)$  arbitrarily. According to (11) we have

$$\begin{aligned} \rho(h, g_n) &\geq \frac{1}{2^i \mu(X_i)} \int_{h^{-1}(M) \cap X_i}^* \frac{|h - g_n|}{1 + |h - g_n|} d\mu^* \geq \\ &\geq \frac{1}{2^i \mu(X_i)} \mu^*(h^{-1}(M) \cap X_i) \cdot \frac{\min_{1 \leq k \leq m_n} r_{nk}}{1 + \min_{1 \leq k \leq m_n} r_{nk}} \geq \frac{1}{2^i \mu(X_i)} \cdot \frac{\mu(X_i)}{j} \cdot \frac{a_M \eta_n}{1 + a_M \eta_n} > \\ &> \frac{1}{2^i j} \cdot \frac{a_M \eta_n}{1 + \eta_n} = \delta_n. \end{aligned}$$

It means by (13) that  $B_\rho(g_n, \delta_n) \subset B_\rho(f_n, r'_n) \setminus \tilde{A}_{i,j}(M)$ . Then in virtue of (9) we get  $B_\rho(g_n, \delta_n) \subset B_\rho(f, r_n) \setminus (F \cup \tilde{A}_{i,j}(M))$ . Consequently

$$\gamma(f, r_n, F \cup \tilde{A}_{i,j}(M)) \geq \delta_n > \frac{a_M \alpha r_n}{6j},$$

which justifies the porosity of  $F \cup \tilde{A}_{i,j}(M)$  at  $f$ .  $\square$

**Theorem 2.** *Let  $M$  be a  $\sigma$ -very porous set. Then  $A(\chi_{\mathbb{R} \setminus M})$  is  $\sigma$ -superporous in  $(\mathcal{M}, \rho)$ .*

*Proof.* We may already suppose that  $M = \bigcup_{k=1}^{\infty} M_k$ , where  $M_k$  is globally very porous and  $a_{M_k} < 1$  for all  $k \in \mathbb{N}$ .

Choose  $f \in A(\chi_{\mathbb{R} \setminus M})$ . Then we have

$$\begin{aligned}
+\infty &= \mu(X) - \int_X |\chi_{\mathbb{R} \setminus M} \circ f| d\mu^* = \mu(X) - \mu^*(f^{-1}(\mathbb{R} \setminus M)) \leq \\
&\leq \mu^*(f^{-1}(M)) \leq \sum_{i,k \in \mathbb{N}} \mu^*(f^{-1}(M_k) \cap X_i),
\end{aligned}$$

thus  $\mu^*(f^{-1}(M_k) \cap X_i) > 0$  for some  $i, k \in \mathbb{N}$ . It suffices now to pick  $j \in \mathbb{N}$  such that  $\mu^*(f^{-1}(M_k) \cap X_i) \geq \frac{\mu(X_i)}{j}$  then clearly  $f \in \tilde{A}_{i,j}(M_k)$ , consequently

$$A(\chi_{\mathbb{R} \setminus M}) \subset \bigcup_{i,j,k \in \mathbb{N}} \tilde{A}_{i,j}(M_k),$$

which concludes the proof by Lemma 4.  $\square$

Now we turn to characterizing the meagerness of  $A(\chi_{\mathbb{R} \setminus M})$  in  $(\mathcal{M}, \rho)$  in terms of properties of  $M$ . We will need the following

**Lemma 5.** *If  $M$  is meager at some point of  $\mathbb{R}$ , then  $A_{i,j}(M)$  is meager at some point of  $(\mathcal{M}_i, \rho_i)$  for all  $i, j \in \mathbb{N}$ .*

*Proof.* In the sequel we will use that the topology induced by  $\rho_i$  on  $\mathcal{M}_i$  is equivalent with the topology of convergence in measure on  $X_i$ , i.e. with the topology induced by the metric  $m_i = m|_{\mathcal{M}_i \times \mathcal{M}_i}$  (see [Ha], Problem 42(4)).

Suppose that there exists an interval  $U = B(t_0, r)$  ( $t_0 \in \mathbb{R}, r > 0$ ) such that  $U \cap M = \bigcup_{k=1}^{\infty} M_k$  for some nowhere dense sets  $M_k \subset \mathbb{R}$  ( $k \in \mathbb{N}$ ). Without loss of generality we may assume that  $M_k \subset M_{k+1}$  for all  $k \in \mathbb{N}$ . Let  $f_0 \equiv t_0$  on  $X_i$  and put  $V = B_{m_i}(f_0, r)$ .

We will show that  $V \cap A_{i,j}(M_k)$  is nowhere dense in  $(\mathcal{M}_i, m_i)$ : take an open ball  $B_{m_i}(f, \varepsilon)$  in  $\mathcal{M}_i$ . We may already suppose that  $f \in V$  and  $f$  equals a simple function  $\sum_{s=1}^m b_s \chi_{D_s}$  where  $b_1, \dots, b_m \in U$  and  $D_1, \dots, D_m$  is a measurable partition of  $X_i$ .

Then the nowhere density of  $M_k$  in  $\mathbb{R}$  yields some  $b'_s \in \mathbb{R}$  and  $0 < \varepsilon_0 < \frac{\mu(X_i)}{j}$  such that

$$(14) \quad B(b'_s, \varepsilon_0) \subset B(b_s, \varepsilon) \setminus M_k \text{ for any } 1 \leq s \leq m.$$

Choose  $g \in B_{m_i}(f_1, \varepsilon_0)$  where  $f_1 = \sum_{s=1}^m b'_s \chi_{D_s}$  then by (14)

$$g^{-1}(M_k) \subset \{x \in X_i; |f_1(x) - g(x)| \geq \varepsilon_0\}.$$

Therefore  $\mu^*(g^{-1}(M_k)) \leq \varepsilon_0 < \frac{\mu(X_i)}{j}$ , so  $g \notin A_{i,j}(M_k)$ . On the other hand  $f_1 \in B_{m_i}(f, \varepsilon)$ ; thus,

$$\emptyset \neq B_{m_i}(f, \varepsilon) \cap B_{m_i}(f_1, \varepsilon_0) \subset B_{m_i}(f, \varepsilon) \setminus A_{i,j}(M_k),$$

which justifies the nowhere density of  $V \cap A_{i,j}(M_k)$  in  $\mathcal{M}_i$ .

Finally, denote  $V_0 = B_{m_i}(f_0, r_0)$  where  $r_0 = \min\{r, \frac{1}{j}\}$ . Pick  $h \in A_{i,j}(M) \cap V_0$ . Then  $h^{-1}(M \setminus U) \subset \{x \in X_i; |h(x) - f_0(x)| \geq r_0\}$ , so  $\mu^*(h^{-1}(M \setminus U)) \leq r_0 \leq \frac{1}{j} < \frac{\mu(X_i)}{2j}$ . Furthermore in view of the regularity of  $\mu^*$  we get (cf. [F], Section 2.1.5(1))

$$\begin{aligned}
\frac{\mu(X_i)}{j} &\leq \mu^*(h^{-1}(M)) \leq \mu^*(h^{-1}(M \cap U)) + \mu^*(h^{-1}(M \setminus U)) < \\
&< \lim_{k \rightarrow \infty} \mu^*(h^{-1}(M_k)) + \frac{\mu(X_i)}{2j},
\end{aligned}$$

hence  $\lim_{k \rightarrow \infty} \mu^*(h^{-1}(M_k)) > \frac{\mu(X_i)}{2^j}$ , so  $h \in A_{i,2j}(M_k) \cap V_0 \subset A_{i,2j}(M_k) \cap V$  for some  $k \in \mathbb{N}$ . Therefore

$$A_{i,j}(M) \cap V_0 \subset \bigcup_{k=1}^{\infty} A_{i,2j}(M_k) \cap V$$

which means that  $A_{i,j}(M)$  is meager at  $f_0$  in  $\mathcal{M}_i$ .

**Theorem 3.** *If  $M$  is meager at some point of  $\mathbb{R}$  then  $A(\chi_{\mathbb{R} \setminus M})$  is meager in  $(\mathcal{M}, \rho)$ .*

*Proof.* Let  $t_0 \in \mathbb{R}$  and  $r > 0$  be such that  $B(t_0, r) \cap M$  is meager in  $\mathbb{R}$ . Let  $V_i = B_{m_i}(f_0, r_0)$  where  $f_0 \equiv t_0$  on  $X$  and  $0 < r_0 = \min\{r, \frac{1}{2}\}$ . Then by Lemma 5  $A_{i,2}(M) \cap V_i$  is meager in  $(\mathcal{M}_i, \rho_i)$  for all  $i \in \mathbb{N}$ .

Choose  $f \in A(\chi_{\mathbb{R} \setminus M})$ . Then  $\mu^*(f^{-1}(\mathbb{R} \setminus M)) < +\infty$  and by the regularity of  $\mu^*$  there exists a  $\mu^*$ -hull  $B$  of  $f^{-1}(\mathbb{R} \setminus M)$  (see [F], Section 2.1.4). Consequently,  $\mu(B \cap X_i) = \mu^*(f^{-1}(\mathbb{R} \setminus M) \cap X_i) = \mu^*(X_i \setminus (X_i \cap f^{-1}(M)))$ ; thus,

$$+\infty > \mu^*(f^{-1}(\mathbb{R} \setminus M)) = \mu(B) = \sum_{i=1}^{\infty} \mu(B \cap X_i) = \sum_{i=1}^{\infty} \mu^*(X_i \setminus (X_i \cap f^{-1}(M))).$$

Then for all  $i \geq m$  ( $m \in \mathbb{N}$ ) we have

$$\frac{\mu(X_i)}{2} > 1 > \mu^*(X_i \setminus (X_i \cap f^{-1}(M))) \geq \mu(X_i) - \mu^*(X_i \cap f^{-1}(M)),$$

hence  $f|_{X_i} \in A_{i,2}(M)$  for all  $i \geq m$ . Accordingly,

$$A(\chi_{\mathbb{R} \setminus M}) \subset \bigcup_{m=1}^{\infty} P_m \text{ where } P_m = \prod_{i=1}^{m-1} \mathcal{M}_i \times \prod_{i=m}^{\infty} A_{i,2}(M) \text{ for each } m \in \mathbb{N}.$$

It suffices now to show by Remark 1 that  $P_m$  is meager in  $P = \prod_{i=1}^{\infty} \mathcal{M}_i$  for every  $m \in \mathbb{N}$ : Let  $\mathbf{U} = \prod_{i=1}^n U_i \times \prod_{i=n+1}^{\infty} \mathcal{M}_i$  be any basic open set of the product topology on  $P$  such that  $n \geq m$ . Denote by  $\mathbf{V}$  the open set  $\prod_{i=1}^n U_i \times V_{n+1} \times \prod_{i=n+2}^{\infty} \mathcal{M}_i \subset P$ . Then  $\mathbf{V} \subset \mathbf{U}$  and  $\mathbf{V} \cap P_m \subset \prod_{i=1}^n U_i \times (V_{n+1} \cap A_{n+1,2}(M)) \times \prod_{i=n+2}^{\infty} A_{i,2}(M)$ , which is meager in  $P$ . It means by Theorem 1.7. in [HMC] that  $P_m$  is meager in  $P$ .  $\square$

**Corollary.**  *$A(\chi_{\mathbb{R} \setminus M})$  is meager in  $(s, \rho_F)$  if and only if  $M$  is meager at some point of  $\mathbb{R}$ .*

*Proof.* The sufficiency follows from the previous theorem by putting  $X = \mathbb{N}$ ,  $S = \mathcal{P}(\mathbb{N})$  and the counting measure on  $\mathbb{N}$  for  $\mu$ .

Conversely, suppose that  $M$  is non-meager everywhere in  $\mathbb{R}$ . Then  $M$  with the relative topology is a dense Baire subspace of  $\mathbb{R}$ . Then the product  $E = M^{\mathbb{N}}$  is a Baire space which is clearly dense in  $s$  ([HMC], Lemma 5.6.). Therefore  $E$  is non-meager in  $s$  and hence  $A(\chi_{\mathbb{R} \setminus M}) \supset E$  is non-meager in  $s$ .  $\square$

*Remark 2.* In connection with the Corollary a question arises if a similar characterization of  $A(\chi_{\mathbb{R} \setminus M})$  is possible also in  $\mathcal{M}$ . Mimicking the above proof and using Remark 1 it would be sufficient to prove that non-meagerness of  $M$  everywhere in  $\mathbb{R}$  implies non-meagerness of  $A_{i,0}$  everywhere in  $\mathcal{M}_i$  for each  $i \in \mathbb{N}$ , further that  $\mathcal{M}_i$  is separable for each  $i \in \mathbb{N}$ . This last condition is needed for the theorem on product of Baire spaces ([HMC], Lemma 5.6.), thus we may consider the question only for separable measure spaces  $(X, S, \mu)$  (see [Ha], §41).



It is not hard to show that this is really the case if each  $X_i$  is a finite disjoint sum of atoms, however in general the answer is not known to me.

*Remark 3.* Another question here arises in connection with finding necessary conditions for  $\sigma$ -porosity of  $A(\chi_{\mathbb{R} \setminus M})$  in  $\mathcal{M}$  (or at least in  $s$ ). If we want to use some argument similar to that of in the Corollary, we would need some ‘‘porosity-Baire’’ product theorem as the mentioned result of Oxtoby ([O1],[HMC]). This ultimately breaks down to proving a porosity version of the well-known Kuratowski-Ulam theorem on sections of nowhere dense subsets of the product space ([O2], Theorem 15.1). More precisely, the questions are as follows:

(i) If  $X$  and  $Y$  are separable metric spaces and  $E$  is a porous subset of  $X \times Y$  with (say) the box metric, then are the  $x$ -sections  $E_x$  of  $E$  porous in  $Y$  except for a  $\sigma$ -porous set in  $X$ ?

(ii) Call a metric space  $Z$   $p$ -Baire if every nonempty open subset of  $Z$  is non- $\sigma$ -porous. Is the property of being separably  $p$ -Baire (countably) productive?

The preceding theorems provide sufficient background for investigating the class

$$\mathcal{U} = \{\Phi \in \mathcal{F}; A(\Phi) \text{ is } \sigma\text{-superporous in } (\mathcal{M}, \rho)\}.$$

**Theorem 4.** *We have*

- (i)  $\text{card}(\mathcal{U} \cap \mathcal{F}_m) = \text{card}\mathcal{U} = 2^c$
- (ii)  $\text{card}(\mathcal{F} \setminus \mathcal{U}) = 2^c$  for  $(s, \rho_F)$ .

*Proof.* (i) Every subset of the Cantor’s ternary set  $C$  is very porous therefore in view of Theorem 2  $\chi_{\mathbb{R} \setminus E} \in \mathcal{U} \cap \mathcal{F}_m$  for every  $E \subset C$ , further  $\chi_{\mathbb{R} \setminus E} \neq \chi_{\mathbb{R} \setminus E'}$  provided  $E \neq E'$ . Consequently  $\text{card}(\mathcal{U} \cap \mathcal{F}_m) \geq \text{card}\mathcal{P}(C) = 2^c$ . Further clearly  $\text{card}\mathcal{U} \leq \text{card}\mathcal{F} \leq \text{card}(\mathbb{R}^{\mathbb{R}}) = 2^c$ .

(ii) If we restrict ourselves to  $(s, \rho_F)$  only, then  $\chi_E \notin \mathcal{U}$  for each  $E \subset C$  since  $A(\chi_E) = s \setminus A(\chi_{\mathbb{R} \setminus E})$  and  $(s, \rho_F)$  is a nonmeager space by Lemma 2. Thus again  $2^c = \text{card}\mathcal{P}(C) \leq \text{card}(\mathcal{F} \setminus \mathcal{U}) \leq \text{card}\mathcal{F} \leq 2^c$ .  $\square$

Further we have

**Theorem 5.**  *$\mathcal{U}$  is residual in  $\mathcal{F}$ .*

*Proof.* See [TZs], Lemma 2 and our Theorem 1.  $\square$

*Remark 4.* It is worth noticing that if we restrict our investigations onto  $\mathcal{F}_m$  only, then similar results hold. Actually, Lemma 3-4 and Theorem 1-2 hold without change, we need only to replace  $\mu^*$  by  $\mu$  and the upper integral by integral, respectively in the proofs.

We can also prove the analogue of Tóth’s Theorem (Theorem 5) for  $\mathcal{F}_m$ :

**Theorem 5’.**  *$\mathcal{U} \cap \mathcal{F}_m$  is residual in  $(\mathcal{F}_m, d)$ .*

*Proof.* See Lemma 2 in [TZs]. The only difference is in proving the density of  $\mathcal{U}_0 = \{\Phi \in \mathcal{F}_m; \Phi \text{ satisfies (6) for some } t_0 \in \mathbb{R}\}$  in  $(\mathcal{F}_m, d)$ , more precisely in proving that  $\Psi = \Phi\chi_M + \frac{\varepsilon}{4}\chi_{\mathbb{R} \setminus M} \in \mathcal{F}_m$ , where  $\Phi \in \mathcal{F}_m, \varepsilon > 0$  and  $M = \{t \in \mathbb{R}; \text{ either } t \notin (0, 1) \text{ or } t \in (0, 1) \text{ and } |\Phi(t)| \geq \frac{\varepsilon}{4}\}$ .

To show this pick  $f \in \mathcal{M}, c \in \mathbb{R}$  arbitrarily and observe that

$$(\Psi \circ f)^{-1}([c, +\infty)) = \begin{cases} (\Phi \circ f)^{-1}([c, +\infty)), & \text{if } c > \frac{\varepsilon}{4} \\ (\Phi \circ f)^{-1}([c, +\infty)) \cup (f^{-1}((0, 1)) \cap (\Phi \circ f)^{-1}((-\frac{\varepsilon}{4}, \frac{\varepsilon}{4}))), & \text{if } c \leq \frac{\varepsilon}{4} \end{cases}$$

thus  $\Psi \circ f \in \mathcal{M}$ .  $\square$

#### REFERENCES

- [AB] S. Agronsky - A. Bruckner, *Local compactness and porosity in metric spaces*, Real Analysis Exchange **11** (1985-86), 365-379.
- [D] E.P. Dolženko, *Boundary properties of arbitrary functions*, Izv. Akad. Nauk SSSR Ser. Mat. **31** (1967), 3-14 (Russian).
- [F] H. Federer, *Geometric Measure Theory*, Springer-Verlag, Berlin-Heidelberg-New York, 1969.
- [G] J.I. Gribanov, *On metrization of a space of functions*, CMUC **4** (1963), 43-46 (Russian).
- [Ha] P.R. Halmos, *Measure Theory*, D. van Nostrand, Toronto-New York-London, 1950.
- [HMC] R.C. Haworth - R.A. McCoy, *Baire spaces*, Dissertationes Math. **141** (1977).
- [K] J.L. Kelley, *General Topology*, Springer-Verlag, New York, 1984.
- [KG] A.A. Kirillov - A.D. Gvishiani, *Theorems and Problems in Functional Analysis*, Springer, New York, 1982.
- [O1] J.C. Oxtoby, *Cartesian products of Baire spaces*, Fund. Math. **49** (1961), 157-166.
- [O2] J.C. Oxtoby, *Measure and Category*, Springer, New York, 1971.
- [Re] D.L. Renfro, *A Study of Porous and  $\sigma$ -Porous Sets*, preprint.
- [TZs] J. Tóth - L. Zsilinszky, *On a typical property of functions*, Math. Slovaca **45** (1995), 121-127.
- [W] W. Wilczyński, *A category analogue of the density topology, approximate continuity and the approximate derivative*, Real Analysis Exchange **10** (1984-85), 241-265.
- [Za1] L. Zajíček, *Porosity and  $\sigma$ -porosity*, Real Analysis Exchange **13** (1987-88), 314-350.
- [Za2] L. Zajíček, *Porosity,  $\mathcal{I}$ -density topology and abstract density topologies*, Real Analysis Exchange **12** (1986-87), 313-326.
- [Za3] L. Zajíček, *Sigma-porous sets in topologically complete metric spaces*, preprint.
- [Zs1] L. Zsilinszky, *On subspaces of measurable real functions*, Portugaliae Math. **52** (1995), 125-130.
- [Zs2] L. Zsilinszky, *On normability of a space of measurable real functions*, Acta Math. et Inf. **2** (1995), 73-76.